



International Journal Of Scientific And University Research Publication

ISSN No **2364/2018**

Listed & Index with
ISSN Directory, Paris



Multi-Subject Journal



THE PREVALENCE OF STAPHYLOCOCCI IN SAUSAGES AT KHARTOUM STATE

Nagwa B.Elhag || Corresponding author: Assistant Professor. Department of Food Hygiene and Safety

College of Public & Environmental Health

University of Bahri

Khartoum North

Sudan

Foods free of Staphylococcal food poisoning organisms have a great concern by consumers nowadays. The aim of this study is to characterize the population of Staphylococci in

staphylococci in sausages obtained from different sources in Khartoum State. A total of 40 fresh sausage samples were collected randomly from (a) butcheries in Khartoum, Khartoum North, and Omdurman, (b) factory processed sausages at retail outlets and (c) homemade sausages obtained from households in Khartoum. Enumeration and characterization of different species of Staphylococci were carried out. The identification of the isolates was mainly based on morphological properties and biochemical tests. Staphylococcal mean counts in all samples ranged from 1.48×10^5 cfu/g to 7.66×10^6 cfu/g. The coagulase positive *Staphylococcus aureus* was identified in all sausage samples that were investigated, making 51.06% of the total isolates, while the coagulase negative isolates were identified as *Staphylococcus epidermidis* (31.91%), *Staphylococcus capitis* (12.77%), *Staphylococcus simulans* (2.13%), and *Staphylococcus hominis* (2.13%). These results indicate that sausage samples are highly contaminated with *Staphylococcus* species particularly *Staphylococcus aureus*. The high incidence of this bacterium in butcheries, processed and home-made sausages is a potential health hazard which indicate improper processing procedure, handling and bad personal habits. Utilization of high quality meat for sausage production and careful handling during processing is necessary to produce sausage free of pathogens.

Sausages, Coagulase positive
staphylococci, Coagulase negative

مقدمة

meat and kofta sold in Khartoum State. As reported by Rabba Mohammed, Mashair, Sulieman and Elgasim (2013) the count (log 4 cfu/g) was recorded for *Staphylococcus aureus* highest prepared Sudanese fresh beef sausages without bee honey, while the lowest count (log 1.5 cfu/g) was recorded for sausages with 7.5% bee honey. Nowadays sausages are becoming an important food item in Sudanese meals. They are usually not fully cooked when sold at retail points as cafeterias and restaurants which make this product a health hazard. The objective of this piece of work is to study the prevalence species in fresh Sudanese sausages sold in Khartoum State.

Collection of samples

A total of 40 fresh sausage samples were collected from eight different sources (5 samples each). These included (a) butcheries in Khartoum, Khartoum North, and Omdurman, (b) Factory processed sausage at retail outlets (F1, F2, F3, and F4), (c) homemade sausages obtained from households in Khartoum. Samples were collected aseptically and kept in sterile insulated ice containers and immediately transferred to the laboratory for analysis.

Microbiological analysis

Thirty grams of each sample of fresh sausages were weighed aseptically, added into sterile conical flask containing 270 ml sterile peptone water and then blended for 30 sec in a sterilized electric blender. Serial tenfold dilutions were prepared following the method described by Harrigan (1998). From suitable dilutions, 0.1 ml was drawn aseptically and spread onto the surface of dried Baird Parker agar medium using sterile glass rod then the plates were incubated at 37°C for 48 hours. Counting of using a colony counter (Quebec Colony Counter) and results were expressed as colony forming unit per gram (cfu/g) of the sample.

Purification and identification of isolates

Typical colonies of *Staphylococcus* were purified and identified following the method described in Bergey's Manual (Sneath, Mair, Sharp & Holt. 1986), Barrow & Felthman (1993) and Harrigan (1998).

Results and Discussion

The pH values and total Staph count are presented in Table 1. The

are highly perishable and are susceptible to contamination by spoilage and pathogenic microorganisms. Different types of microorganisms may be introduced into sausages from various sources due to improper storage temperature and manipulation and poor hygienic condition in non industrial establishments especially in butcheries (Cordoba, Cordoba & Jordano, 1998). Some of these microorganisms can cause food poisoning and infections when they are introduced in high numbers due to direct contact with food handlers (Byrant, Javis & Gilbet, 1988). Children and elderly people are the more susceptible group to food infections (Patricia Blas & Ana, 2006). Staphylococcal food poisoning is gastrointestinal illness caused by food contaminated is normally found *Staphylococcus aureus*. *Staphylococcus aureus* with in different parts of human beings and other animals as in skin, the mucosal membrane, mouth, nose, pimples, boils, different environments, and in a wide range of food stuff (Acco, Ferreira, is usually present *Staphylococcus aureus* Henriques, & Tondo, 2003). in small numbers in raw meat and in food handlers. Human beings, different types of foods and environment are considered as favorable transmission (Lateef, 2004). *Staphylococcus aureus* environment for Angela Cheryl and Charles (2005) reported that up to 25% of healthy in their skin and in nose. Keeping *Staphylococcus aureus* people have of meat and meat products free from contamination species is difficult, therefore the food processor *Staphylococcus* with should keep meat and meat products at temperature that prevent the spp. Meat and meat products safety *Staphylococcus* growth of depends on quality of raw materials used, their adequate handling, processing, preparation, transportation, marketing, and storage. Generally staphylococcal food poisoning strains come from human sources and 95% of food poisoning outbreaks are caused by enterotoxin A, B, C, D and E (Letetre, Perelle, Dilasser & Fach 2003). The typical symptoms of food poisoning required very low amount of enterotoxin ranging from 20 µg to 1 µg (US Food and Drug Administration, 2001) which corresponding to 105 Staphylococcal colony forming units per gram of foods (Normanno Salandra, Dambrosio, Quaglia, Corrente, Parisi, Santagada, Firinu, Crisetti & Celano, 2007). Nadia (2010) found that sausages have the with the rate of 72% when *Staphylococcus aureus* highest count of compared with the other Sudanese meat products (burger, minced

in fresh sausages samples stored at 4°C for 10 days (Rantsiou, Iacumin, Urso, Cantoni, Comi and Coccolin, 2005). They found that *S. aureus* (50%), strains of *Staphylococcus xylophilus* most of the isolates were, were also identified. *S. succinus* and *Staphylococcus* *S. warneri* *pasteuri* Coccolin, Rantsiou, Iacumin, Urso, Cantoni, and Comi, (2004) *Staphylococcus xylophilus* was isolated from fresh sausages reported that nowadays *S. epidermidis* 4°C for 10 days. The identified *S. aureus* stored at seen as an opportunistic pathogen. It is now the most frequent cause of nosocomial infections at a rate about as high as that due to its. The fact that they are extremely *S. aureus* more virulence cause difficult to treat represent a serious burden for the public health is part of normal flora of the skin of *capitis* *S. aureus* (Otto, 2009). the human scalp, face, neck and ears and has been associated with prosthetic valve of endocarditis (D'mello Daley, Rahman, Qu, is *S. simulans* Garland, Pearce and Deighton, 2008). The identified frequently acquired through contact with domestic animals. It is an authentic pathogenic agent of osteoarticular infection (Mallet, Loiez, may *hominis* *S. aureus* Melliez, Yazdanpanah, Senneville, Lemaire, 2011). occasionally cause infection in patients whose immune system are compromised for example by chemotherapy (Kloos & Schleifer, 1975). Poor hygiene practices such as negligence to wash hands after visiting the bath room may result in up to 107 pathogens under the fingernails of the handlers (Nel Lues, Buys and Venter, 1993). The and the presence of *Staphylococcus aureus* high prevalence rate of coagulase negative staphylococci in the sausage samples under study indicates the poor hygienic conditions during sausages production in Sudan. Also it reflects the poor personal hygiene of people dealing with food processing. Lack of standards hygiene measures among workers in factories processing sausage may lead serious health hazards to consumers. These type of research make a critical step towards understanding the serious prevalence of staphylococci in sausage produced under poor processing conditions.

استنتاج

is frequently acquired through *S. simulans* The identified contact with domestic animals. It is an authentic pathogenic agent of osteoarticular infection (Mallet, Loiez, Melliez, may *hominis* *S. aureus* Yazdanpanah, Senneville, Lemaire, 2011). occasionally cause infection in patients whose immune system are compromised for example by chemotherapy (Kloos & Schleifer, 1975). Poor hygiene practices such as negligence to wash hands after visiting the bath room may result in up to 107 pathogens under the fingernails of the handlers (Nel Lues, Buys and Venter, 1993). The high prevalence rate and the presence of coagulase *Staphylococcus aureus* of negative staphylococci in the sausage samples under study indicates the poor hygienic conditions during sausages production in Sudan. Also it reflects the poor personal hygiene of people dealing with food processing. Lack of standards hygiene measures among workers in factories processing sausage may lead serious health hazards to consumers. These type of research make a critical step towards understanding the serious prevalence of staphylococci in sausage produced under poor processing conditions.

ref_str

- . M. & Mohammed, M. T. (1989). Beef casing and finished **Abbar, F. I.** beef sausages as a source of Salmonella in Iraq. Journal of Food Protection. 52:254-255. I
.; Ferreira, F. S., Henriques, J. A. P., & Tondo, E. C. (2003). **Acco, M. J.** Identification of multiple strains of *Staphylococcus aureus* colonizing nasal mucosa of food handlers. Food Microbiology. 20:

mean pH values of the sausage samples ranged between 5.83 and 6.23. The highest mean pH values were recorded in samples. (2004) reported that the pH *in vacuo* obtained from butcheries. values for fresh sausages is not lower than 5.5. Results showed that the pH of most samples investigated were higher than this value. The high mean pH values for the sampled sausages revealed that these sausages were produced from low quality meat with pH values higher than 6.00, which is referred to as dark, firm and dry meat (DFD). Newton and Gill (1981) claimed that DFD meat is characterized by high ultimate pH > 6.00 and deficiencies in glucose because of exercise or stress. These factors can result in bacterial spoilage becoming evident at an earlier stage of growth of the meat flora. Newton and Gill (1981) found that the pH value is an important indicator for microbial growth and high pH resulted in high microbial count and short shelf life. The ultimate pH of meat is significant for its resistance to spoilage because most bacteria grow optimally at about pH 7.0 and not below pH 4.0 or above pH 9 (Jamilah, Abbas, and Abdoul 2008). Staphylococcal mean counts ranged from 1.48x10⁵ to 7.66x10⁶cfu/g. Factory 2 recorded the highest count (7.66x10⁶cfu/g), while the lowest mean count was reported for samples obtained from factory 3 (Table 1). Similar results were obtained by Oluwafemi and Simisaye (2006) who found counts ranged from 1.30x10⁵cfu/g to *S. aureus* that the 2.20x10⁷cfu/g for sausage samples obtained from markets in Nigeria. The mean values of Staphylococcal counts of the investigated sausage samples were higher than the standard limits (5x10²cells) established by the Sudanese Microbiological Standards for Foods (SSMO, 2001). The number of these bacteria in sausages is usually low as obtained in Sudan by Ali (2004), who found count of sausages at zero day of storage at *Staphylococcus* that refrigerator temperature was 1.17x10²cfu/g, reached 1.19x10²cfu/g after 5 days of storage and 2.8x10²cfu/g at the end of 10 days of (2011) recorded a lower storage. Samapito, Leenanon, and Robert count of staphylococci (3.3x10³-2.7x10⁶cfu/g) for "Mhom", a traditional meat sausage in Thailand, than that obtained in this study. The high counts of this bacteria in the sampled sausages may be attributed to crosscontamination during preparation, processing, transportation and packaging as has been observed by Abbar & Mohammed (1989). In addition, some bad habits of butchers such as wiping and dusting of displayed sausage with dirty pieces of cloth that are used for cleaning hands and tables, sneezing and snuffing play an important source of contamination. Humans are the principal source or reservoir of these organisms. Staphylococci are found in the nose, throat and skin of up to 60% of healthy humans. Staphylococcal food poisoning is a persistent cause of gastroenteritis worldwide, especially in the developing countries (Vora, Senecal, and Schaffner, 2003). Concerning public health Staphylococci are important organisms that may occur in cooked comminuted meat products. They are primarily found in processed meat and dairy products; survive in the salted medium of hams and sausages (Jay, sp. are presented *Staphylococcus* 1996). The identified isolates of the *Staphylococcus* in Table 2. Results showed that the coagulase positive was identified in all sausage samples that were investigated, *aureus* making 51.06% of the total isolates. Most of the isolates from Khartoum, Khartoum North and Omdurman butcheries were *Staphylococcus aureus* (25.53%). *Staphylococcus aureus* identified as is considered the third most important cause of disease in the world amongst the reported foodborne illnesses (Anonou, Maqueda, Martine, Bueno, Galvez and Valdivia, 2007). It is a saprophyte and commensal of the skin mucus membrane in both animal and humans. As a pathogen it can cause a number of diseases ranging from minor *S. aureus* skin infection to fetal sepsis. Toxigenic strains of currently among the leading causes of food borne intoxication. The *Staphylococcus* coagulase negative staphylococci were identified as (2.13%), *S. simulans* (12.77%), *S. capitis* (31.91%), *epidermidis* (2.13%). Presence of staphylococci in Sudanese *S. hominis* and spp. *Staphylococcus* sausages is not unique as other authors identified

- of Biomedical Research. 9 (2): 133-136.
- (2009). Staphylococcus epidermidis the "accidental" pathogen. *Nature Review Microbiology*. 7: 555-567. doi:10.1038/nrmicro2182.
- , Aicalizzi, B., & Ana, M. S. (2006). Prevalence of Some Bacteria, Yeasts and Molds in Meat Food in San Luis, Argentina. *Cent Eur. J. Publ. Health*. 4(3): <http://apps.szu.cz/svi/ceiph/archiv/2006-3-10-full.pdf>, 141-144.
- Mohammed, Mashair A. Sulieman & Elgasim A. E. (2013). Effect of Bee Honey in Safety and Storability of Beef Sausage. *Pakistan Journal of Nutrition* 12 (6): 560-566. doi:10.3923/pjn.2013.560.566.
- ., Iacumin, L., Urso, R., Cantoni, C., Comi, G. & Cocolin, L. (2005). Ecology and characterization by molecular methods of Staphylococcus species isolated from fresh sausages. *Int J Food Microbiol*. 97(3):277-84. <http://www.sciencedirect.com/science/article/pii/S0168160504002661>.
- ., Leenanon, B., & Robert, E. L. (2011). Microbiological characteristics of "Mhom", a Thi traditional meat sausage. *The Open Food Science Journal*. 5: TOFSJ-5-31. <http://benthamopen.com/contents/pdf/TOFSJ/31-36>.
- , Mair, N. S., Sharp, M. E., & Holt, J. G. (eds) (1986). *Berg's Manual of Systematic Bacteriology*. Vol. 2, (Baltimore: Williams and Wilkins).
- (2001). Sudanese Standards and Metrology Organization. Sudanese Microbiological Standards for Foods. Khartoum, Sudan.
- . (2001). Food borne pathogenic microorganisms and Natural **U.S. Food and Drug Administration**. <http://www.cfsan.fda.gov> Toxins.
- , Senecal, A. & Schaffner, D.W. (2003). Survival of *S. aureus* ATCC 13565 in intermediate moisture foods is highly variable. *Risk Anal*. 23(1): 229- 236. DOI: 10.1111/1539-6924.00302
- 489-493.
- <http://www.sciencedirect.com/science/article/pii/S074000200300049>
- . (2004). The Effect of Chilling and Storage on Bacterial Contamination and Quality of Red Meat and its Manufactured Products. M. Sc. Thesis, Faculty of Veterinary Medicine, University of Khartoum, Sudan.
- ., Cheryl, B. O. & Charles, H. (2005). Staphylococcal Food Poisoning Outbreak in Southeast Kansas. Kansas Department. Health Environment (KD,HE). Foodborne illness outbreak report. http://www.kdheks.gov/epi/download/foodborne_illness_outbreak_report_dec_2005.pdf. http://www.kdheks.gov/epi/download/foodborne_illness_outbreak_report_dec_2005.pdf.
- ., Maqueda, M., Martinez Bueno, M., Galvez, A., & Anonou, S. Valdivia, E. (2007). Bactericidal synergism through enterocin AS-48 and chemical preservatives against *S aureus*. *Letters in Appl Microbiol*. 48 (1): 19- 23. <http://www.biolonline.org.br/pdf%3Fmd06023>.
- , & Feltham, R.K.A. (1993). Manual for Identification of Medical Bacteria, third edition. Cambridge University Press, England.
- , Jarvis, J., & Gilbert, G. (1988). Selective enterotoxin production by a Staphylococcus aureus strain indicated in food-borne outbreak. *J. Food Prot*. 51: 130-131.
- ., Rantsiou, K., Iacumin, L., Urso, R., Cantoni, C & Comi, G. (2004). Study of the Ecology of Fresh Sausages and Characterization of Populations of Lactic Acid Bacteria by Molecular Methods. *Appl. Environ. Microbiol*. Vol. 70 (4): 1883-1894. doi: 10.1128/AEM.70.4.1883-1894.2004.
- , Cordoba, J. J., & Jordano. R. (1998). Evaluation of microbial hazards during processing of Spanish prepared Flamerquin. *J. Food. Prot*. 61(6): 693-699.
- , Daley, A.J, Rahman, M.S., Qu, Y., Garland S.; Pearce, ChD'mello D. & Deighton, MA. (2008). "Vancomycin heteroresistance in bloodstream isolates of Staphylococcus capitis". *J. Clin. Microbiol*. 46 (9): 3124–3126. doi:10.1128/JCM.00592-08. PMC 2546727. PMID 18596138.
- . (1998). Laboratory Methods in Food and Dairy Microbiology. Academic press. USA.
- , Abbas, K.A., & Abdoul, R. R. (2008). A Review on some Organic Acids Additives as Shelf Life Extenders of Fresh Beef Cuts. *American. J. Agric. Biol. Sci*.3: 566-574.
- (1996). Fresh meats and poultry. In: *Modern Food Microbiology*. 5th edition. International Thomas Publishing, Kentucky. 63-91.
- , Schleifer K H (1975). Isolation and characterization of staphylococci from human skin. Descriptions of four new sp. *Staphylococcus warneri* *Staphylococcus capitis*, *Staphylococcus hominis* and *Staphylococcus simulans* *International Journal of Systematic Bacteriology* 2562-79. doi: 10.1099/00207713-25-1-62 IJSEM.
- . (2004). The microbiology of pharmaceutical effluent and its public health implications. *World. J. Microbiol. Biotechnol*. 20: 167-171. doi: 10.1023/B:WIBI.0000021752.29468.4e.
- ., Perelle, S., Dilasser, F., & Fach, P. (2003). Identification of new putative enterotoxin SEU encoded by the egc cluster of *Staphylococcus aureus*. *J. Appl. Microbiol*. 95: <http://www.ncbi.nlm.nih.gov/pubmed/12807452> 38-43.
- ., Loiez, C., Melliez, H., Yazdanpanah, Y., Senneville, E., & Lemaire, X., (2011). *Staphylococcus simulans* as an authentic pathogenic agent of osteoarticular infections. *39(5):473-6*. doi: 10.1007/s15010-011-0173-x.
- ., Lues, J., F., R., Buys, E. M. & Venter, P. (1993). The personal and general hygiene practices in the deboning room of a high throughput red meat abattoir. *Food Control* 15: 571-578.
- <http://www.sciencedirect.com/science/article/pii/S095671350300161>
- . G., & Gill, C. O. (1981). The microbiology of dark firm and dry fresh Meats: A Rev. *J. Meat Sci*. 95: 223-232. doi: 10.1016/0309-1740(81)90005-X PMID: 22056031.
- Salandra, L. G., Dambrosio, A., Quaglia, N.C., Corrente, M., Parisi, A., Santagada, G., Firinu, A., Crisetti, E., & Celano, G.V. (2007). Occurrence, characterization and antimicrobial resistance of enterotoxigenic *Staphylococcus aureus* isolated from meat and dairy products. *Int. J. Food Microbiol*. 115: <http://www.ncbi.nlm.nih.gov/pubmed/1732162290-296>.
- , & Simisaye, M.T. (2006). Extent of microbial contamination of sausages sold in two Nigerian cites. *African Journal*



IJSURP Publishing Academy

International Journal Of Scientific And University Research Publication
Multi-Subject Journal

Editor.

International Journal Of Scientific And University Research Publication



+965 99549511



+90 5374545296



+961 03236496



+44 (0)203 197 6676

www.ijsurp.com