



International Journal Of Scientific And University Research Publication

ISSN No **2364/2018**

Listed & Index with
ISSN Directory, Paris



Multi-Subject Journal



BIOPROSPECTING OF BIOCHEMICAL PARAMETERS OF WHEAT UNDER SALINITY STRESS AFTER BIOFERTILIZER TREATMENT

**Prabhat Nath Jha || Department of Biological Science
Birla Institute of Technology and Science (BITS)
Pilani
333031
Rajasthan
India.**

Microbial inoculums termed as 'Biofertilizers' are gaining a lot of interest now a day for improving the growth and productivity as a alternative to chemical fertilizers

and pesticides. In developing countries like India, agriculture is an important sector for economy. However increasing soil salinity causes a major loss of agricultural economy. The present study demonstrated the efficacy of biofertilizer *Enterobacter cloacae* to growth improvement by analyzing the various biochemical parameters of wheat under various level of salinity stress. Microbial isolate was isolated from salt rich lake 'Sambhar lake' to growth improvement and development as biofertilizers for plants growing around the salt rich region. The seeds of wheat were treated with microbial inoculums with salt stress and the observed changes in biochemical parameters were analyzed after 15 days of study period. Results illustrated that plants treated with *Enterobacter cloacae* showed significant improvement in various biochemical parameters responsible for plants sustainability as compared to untreated control plants.

Biofertilizer, Plant growth promoting bacteria, Salt stress, Compatible

مقدمة

with an overall average loss of 40%. Wheat is grown in India under subtropical environments during mild winter which warms up towards grain filling stages of the crop. Wheat is one of the most important cultivated cereals second to rice in importance as staple food in India. However in different parts of India, productivity is affected by increase in salinity as it lowers osmotic potential leading to decreased water availability. Therefore we may speculate that development of microbial inoculants as biofertilizers from extreme condition could be beneficial to mitigate the salt stress to the plants growing in saline habitat. While a number of studies have demonstrated the role of microbes in maintaining the plant growth under unfavourable conditions, little is known about the inoculation on the biochemical parameters of wheat *Enterobacter cloacae* of under different level of salinity. Therefore the aim of the experiment was to increase the productivity of crops with the significant changes in various biochemical properties with the help of biofertilizers.

Materials and methods Bacterial inoculum and seed treatment-1

Invitro pot study was conducted to evaluate the potential of microbial inoculums as biofertilizer. Sterilization of soil was done by autoclaving at 121°C for 1 h for three consecutive days to kill the entire microorganism and their spores. The surface sterilized wheat and kept *Enterobacter cloacae* seeds were treated with biofertilizer for germination in plant growth chamber. After sprouting, plantlets were treated with salt stress 150mM, 175mM,200mM. Bacterized seeds were sown in plastic pots filled with soil (300g per pot) in triplicates in a growth chamber with 16:8 photoperiods up to 21 days at 28 ± 2°C. Control seeds without any treatment were also used in the study. Hoagland media supplemented with salt (150mM, 175mM, 200mM) was used for providing the nutrient as well as imposing the salt treatments. After 15 days of study period, the biochemical parameters such as total proline content, malondialdehyde content, level of total soluble sugar and total protein content were studied by standard methods.

2-Results and discussion

showed *Enterobacter cloacae* Wheat seed treated with biofertilizer significant improvement in biochemical parameters as compared to control plants. Plants treated with biofertilizer showed enhancement of proline, total soluble sugar, total protein content and decrease in malondialdehyde. Our results are agreed with the previous finding of Ruffino et al. (2010) and Nabti et al. (2010).

3-Research Paper

Salinity is one of the major abiotic stresses adversely affecting modern agriculture and constitutes a major problem worldwide. It is estimated that more than 6% of the world's total land area is salt affected (Bui, 2013). Increasing salinity in the soil make it harder for plants root to extract water and therefore have an immediate effect on cell growth and associated metabolism. Accumulation of salinity induced toxic ions affects the plant functionality in several ways. In addition, several biochemical and physiological processes like protein synthesis, lipid metabolism, photosynthesis, ionic homeostasis and nitrogen fixation are severely affected by salinity. One approach to solve the salt stress problem is the use of plant growth-promoting bacteria as biofertilizers. Use of microbial inoculums or biofertilizers is of great agricultural importance as they are cost effective and renewable source of energy for plants and also helps in reducing the use of chemical fertilizers for sustainable agriculture (Rana et al., 2013). Many of the plant growth-promoting bacteria have been reported to colonize the plant rhizosphere and confer beneficial effects by various direct and indirect mechanisms. Term plant growth promoting bacteria was first introduced in 1978 by Kloepper and colleagues to a group of beneficial bacteria that promote the plant by enhancing the acquisition of nutrients, production of phytohormones, nitrogen fixation, siderophore production and indirectly by antibiotic production, secretion of lytic enzymes. The effectiveness of these plant growth promoting bacteria as biofertilizers to mitigate the adverse effect of salinity stress has been reported in several vegetable and other crop plants (Saravankumar and Samiyappan 2007). To maintain osmotic adjustment under salt stress, plants have developed complex physiological and biochemical mechanisms to accumulate the compatible solutes (osmolytes) for maintaining the intracellular osmotic homeostasis (Gupta and Huang 2014). One of the important osmolytes, proline contribute to osmotic adjustment by detoxifying ROS induced oxidative damage (Gupta and Huang 2014). Following inoculation of beneficial bacteria enhanced production of proline (Ait Barka et al. 2006) was observed in abiotically stressed plants. Besides of proline other molecules like glycine betaine, total soluble sugars and total protein content also helps to survival of plants under salinity stress. Previous reports suggested that inoculation with biofertilizers enhance the salt tolerance in tomato (Mayak et al. 2004), red pepper (Siddikee et al. 2011) and groundnut (Saravankumar and Samiyappan 2007). Salt induced land degradation led to a major drop in wheat grain yield ranged 20-43%

tomato plants to salt stress. *Plant Physiol Biochem* 2004; 42:565–572.

Fischer, D., Hofmann, A., Nabti, E., Sahnoune, M., Ghoul, M., Rothballer, M., Schmid, M., Hartmann, A. Restoration of growth of durum wheat (*Triticum durum* var. waha) under saline conditions due to inoculation with the rhizosphere bacterium *Azospirillum brasilense* NH and extracts of the marine alga *Ulva lactuca*. *J Plant Growth Regul* 2010; 29: 6–22.

and Their Role in **Rana R R and Kapoor P Biofertilizers**. *Agriculture; Popular Kheti*, 2013; 1(1):5661.

and Prado, F. The **Ruffino, A., Rosa, M., Hilal, M., González, J.,** role of cotyledon metabolism in the establishment of quinoa (*Chenopodium quinoa*) seedlings growing under salinity. *Plant Soil* 2010; 326: 213–224.

R. ACC deaminase from **Saravanakumar, D., Samiyap- pan,** *Pseudomonas Xuorescens* mediated saline resistance in groundnut (*Arachis hypogea*) plants. *J Appl Microbiol* 2007; 102:1283–1292.
 Chung JB, Sa T Influence of **Siddikee MA, Tipayno SC, Kim K,** varying degree of salinity/sodicity stress on enzyme activities and bacterial populations of coastal soils of Yellow Sea, South Korea. *J Microbiol Biotechnol* 2011; 21:341–346.

on proline *Enterobacter cloacae* Table 1: Effects of biofertilizer

Proline (µmol/g FW)	enttmaerT	content
		alt (0)S
3.17 ± 0.24		UN-INC
4.90±0.18		SL-12

on total *Enterobacter cloacae* Table 2: Effects of biofertilizer

g/g FW)m(SS T	enttmaerT	soluble sugar
		alt (0)S
4.18 ± 0.17		UNINC
5.47±0.19		L12S

استنتاج

Wheat seeds treated with biofertilizer *Enterobacter cloacae* showed significant increase in various biochemical parameters under salinity stress. The specific resistance mechanism of compatible solutes to a wide range of abiotic stresses is still not fully understood, however these molecules significantly contribute to osmotic adjustment, stabilize lipid membranes, stabilising protein during stresses etc. Therefore, use of bacteria with such features could be used as biofertilizer to combat saline stress.

ref_str

Clement C Enhancement of chilling resistance **Ait Barka E, Nowak J,** .1 of inoculated grapevine plantlets with a plant growth-promoting rhizobacterium, *Burkholderia phytofirmans* strain PsJN. *Appl Environ Microbiol* 2006, 72:7246–7252.
 : a neglected factor in plant ecology and **Bui EN. Soil salinity** .2 biogeography. *J Arid Environ.* 2013; 92: 14–25.
 Mechanism of salinity tolerance in plants: **Gupta, B. and Huang, B.** .3 physiological, biochemical, and molecular characteriza- tion. *Int J Genom* 2014;701596. doi:10.1155/2014/701596.
 et al. Plant growth-promoting bacteria confer resistance in **Mayak S.** .4



IJSURP Publishing Academy

International Journal Of Scientific And University Research Publication

Multi-Subject Journal

Editor.

International Journal Of Scientific And University Research Publication



+965 99549511



+90 5374545296



+961 03236496



+44 (0)203 197 6676

www.ijsurp.com