



International Journal Of Scientific And University Research Publication

ISSN No **301/704**

Listed & Index with
ISSN Directory, Paris



Multi-Subject Journal



BIO CONTROL POTENTIAL OF RICE ENDOPHYTE FUSARIUM OXYSPORUM AGAINST RICE SHEATH BLIGHT PATHOGEN RHIZOCTONIA SOLANI

Divya R

Aju K Asok and Jisha MS || Corresponding author: Jisha MS

Associate Professor

School of Biosciences

Mahatma Gandhi University

India.

Discovering novel and effective microbial strains for increasing plant productivity; by enhancing the availability of utilizable minerals and by inhibiting an array of bacterial

and fungal pathogens to plant is innovative approach in sustainable agriculture. There is particular interest in growing crops with little or no dependence on chemical pesticides and placing considerable weight on biological control. Fungal endophytes are a group of microorganisms associated with plants and have attracted a great interest to botanist, microbiologist and ecologist. In this study endophytic fungi were isolated from rice and were evaluated for their antagonism against *Rhizoctonia solani*, an important rice pathogen causing Sheath Blight disease in rice.

Endophytic microorganismالكلمات الرئيسية

Graminaceous crops; *Oryza sativa*;
Fungal

pathogen causing Sheath blight disease in paddy.

مقدمة

Materials and Methods

Rice Plant Samples

Rice plants were collected from Kottayam, Alapuzha and Trivandrum districts, Kerala. Ten rice plants without signs of any disease were collected from four plots of one experimental field.

Samples should be air dried to remove any surface moisture before transport or storage. During transport, it should be kept cool and dry. Paper bags were used. Samples were processed as quickly as possible following collection usually within 24 hours.

Isolated Pathogen and Culture Condition

isolated from infected rice was provided by Rice *hizoctonia solani*, R Research Institute, Mancombu, Kerala. The cultures were frequently sub-cultured and maintained in Potato Dextrose Agar (PDA) medium.

Sample preparation

A vigorous prewashing of plant material was performed to reduce the number of surface microflora and soil. This increases the effectiveness of the surface sterilizer. Prewash was done using running tap water. Samples were surface sterilized using NaOCl solution for 5 minutes [6]. Finally they were rinsed three times in sterile distilled water. The surface sterilized samples were cut into 1cm fragments. The flow out after washing the surface sterilized samples with sterilized water was plated on PDA plate and incubated at room temperature. A control plate with medium was also incubated along with it.

Isolation of endophytic fungi

The surface-sterilized samples were cut into 1cm fragments. Ten fragments from different parts of leaves per rice sample were placed on YEG agar medium. These plates were incubated at 25 °C for 8-12 days. Hyphal tips of the developing fungal colonies were transferred onto PDA. After purifying the isolates several times, the final pure cultures were transferred to PDA slant tubes.

Identification of fungal endophytes

Isolates from slants were transferred to PDA medium on a sterile microscopic slide, which was then covered with a sterile cover glass, incubated at 25 °C and periodically the fungal fruiting structures were observed by microscope using lactophenol cotton blue staining. Further identification of fungal endophyte was achieved by

An endophyte is a fungal or bacterial microorganism that spends the whole or part of its life cycle colonizing healthy tissues of the host plant, typically causing no apparent symptoms of disease [1]. The relationship between the host and the endophyte may range from symbiotic to near pathogenic [2]. Endophytic microorganisms are a significant reservoir of genetic diversity, and an important source for the discovery of novel bioactive secondary metabolites. Endophytes are a rich source of natural products displaying a broad spectrum of biological activities and the phytochemistry of endophytic microbes continues to increase in significance. As a general rule, a single endophytic strain will produce multiple bioactive substances. The reported natural products from endophytes include antibiotics, antipathogens, immunosuppressants, anticancer compounds, antioxidant agents and other biologically active substances. Many medicinal plants are reported to be harboring endophytic fungi.

The ecological roles played by endophytic fungi are diverse and varied [3]. Besides protection against insects, many of these fungi produce biologically active secondary metabolites [4]. Certain, other metabolites produce host growth responses, accelerate or delay senescence [3]. Future investigations might include studies aimed at detecting production of antibiotics and pest deterrents in plants as a first step towards evaluating ecological significance of secondary metabolite production by endophytes in biological control.

, one of the primary graminaceous crops, (*Oryza sativa*) Rice constitutes the main nutrient resource for 40% of world's population including most developing countries. Rice has been severely affected by several diseases and insect pests, of which Sheath blight (ShB) is becoming a serious concern *Rhizoctonia solani* disease caused by for the successful cultivation of rice. It is one of the most destructive diseases of rice that occurs globally [5]. Abuse of chemical pesticides, which are the most common approach for control, can destroy the balance of eco systems and the contamination by their toxic residues may cause harm to humans and domestic animals. In addition, chemicals can decrease the soil micro flora sharply and induce tolerance of the pathogens. Search for environmentally and toxicologically safe, more selective and efficacious fungicides for the control of plant diseases is gaining momentum at present. As an environmentally benign agent, endophytes are the ideal candidates. The aim of the present study is which (*Oryza sativa*) to isolate potential endophytic fungi from paddy , an important rice *Rhizoctonia solani* will be effective against

present study describes a novel potential endophyte isolated (*L.*) that appears to be quite effective *Oryza sativa* from rice (and contributes enhanced *Rhizoctonia solani* in inhibiting productivity.

Molecular analysis of fungal rDNA at the sequence level provides a powerful technique for assessing fungal diversity at the genus level. The molecular analysis of the isolated endophytic fungus with antagonistic property during the study

Fusarium oxysporum. found that the endophyte is Novelty of the endophytic association is its ability to colonize different host plants and tissues. It was found that it can control the Sheath blight disease in rice. In the preliminary studies, [7], reported that exploitation of the endophytes as biocontrol agents may be a breakthrough in the biological control adding strength to additional benefits of host/fungus endophytic association [8].

ref_str

- (2002) Microbial gifts from rain forests. *Can J Plant* Strobel GA. 1. 24: 14-24. *Pathol.*
- (2000) Endophytic communities of rhizobacteria Sturz AV, Nowak J. 2. and the strategies required to create yield enhancing associations with crops. *Appl. Soil Ecol* 15: 183-190.
- (1998) Fungal Saikkonen K, Faeth SH, Helander M, Sullivan TJ. 3. endophytes: a continuum of interactions with host plants. *Ann Rev* 29: 319-343. *Ecol.*
- (1998) Fungal endophytes of tree Pelaez F, Collado J, Arenal F. 4. pp: 179-197.. leaves. *Microbial ecology*
- Slaton NA, Cartwright RD, Meng J, Gbur EE Jr, Norman RJ. 5. Blight severity and rice yield as affected by nitrogen (2003) Sheath application method and fungicide. *Agronomy Journal* fertilizer rate. 1496. 95(6): 1489-1496.
- (1993) Endophytes from herbaceous Schulz BU, Draeger S, Aust HJ. 6. plants and shrubs: Effectiveness of surface sterilization methods. *Mycol Res* 97: 1447.
- (2008) The Mathivanan N, Prabavathi VR, Vijayanandraj VR. 7. fungal secondary metabolites on bacterial and fungal effect of Petr Karlovsky (Ed.). *Secondary metabolites in soil pathogens*. In: 140. *ecology*. pp. 129-140.
- (1997) Host specific Vindhyasekharan P, Ruby P, Samiyappan R. 8. the rice sheath blight pathogen. *R solani* production by toxin

amplification of Internal Transcribed Spacer regions (ITS) in conjunction with 18S rDNA sequencing. The DNA was isolated and ITS region was PCR amplified with ITS1 and ITS 2 primers.

Amplicon was electrophoresed in a 1% Agarose gel and visualized under UV. Concentration of the amplicon was checked in a Nanodrop ND 2000. The amplicon was purified using Nucleospin purification column (Macherey-Nagel). Sequencing of amplicon with forward and reverse ITS primers in ABI 3730xl cycle sequencer. Forward and reverse sequences were assembled and contig was generated after trimming the low quality bases. The sequence analysis was carried out using bioinformatic tool BLAST of NCBI. Based on maximum identity score first few sequences were selected and aligned using multiple sequence alignment software Clustal W. Dendrogram was constructed using neighbor joining method.

of antagonistic potential of the endophytes aluation veIn vitro (Dual plate assay)

Isolated endophytic fungi were screened for their antagonistic ability (Sheath Blight *Rhizoctonia solani* against an important plant pathogen, of rice) by the dual culture technique. Petri dishes with the growth medium were simultaneously inoculated with 9mm disc of the pathogen as well as antagonist near the periphery at diametrically opposite points. Periodic observations are made at regular intervals on the linear growth (mm) of the antagonist colonizing the pathogen's growth.

استنتاج

This study was done to isolate endophytic fungus from rice *Rhizoctonia* and was evaluated for their antagonism against , an important rice pathogen causing Sheath Blight *solani* disease in rice. Endophytic fungi were isolated from the plant materials after eight to twelve days of incubation. The identification of the endophytic fungi was done by colour, texture, pigmentation and morphology of the colonies and the Lactophenol cotton blue staining of the conidiophores.

The colony of first isolate was wooly and eventually covered by grayish, short, aerial hyphae. The conidiophores were mostly unbranched. The conidia were ellipsoidal with short cylindrical beak with muriform septation. This isolate was sp. The colony of the second isolate *Alternaria* identified as was wooly and black in colour. The conidia was 4-5 cells separated by transverse septa formed from twisted sp. *Curvularia* conodiophores, the characteristic feature of The third isolate produced black colour colony with vesicle bearing conidia. Conidia were in single or paired resemble yeast cells, chains or detached and dispersed. This isolate was sp. The colony of the fourth isolate *Aspergillus* identified as were cottony pinkish white. Conidia were thin walled, three to five septate, fusoid and pointed at both ends. This isolate was sp. *usarium* Fidentified as

Isolated endophytic fungi were tested for antagonistic activity sp. *Fusarium* by dual culture method. The isolated and *Rhizoctonia solani* significantly inhibited the growth of the percentage of inhibition was found to be 90.5%. The level against test *Fusarium oxysporum* of antagonism exhibited by was very high. It showed strong *Rhizoctonia solani* pathogen inhibition of mycelia growth of plant pathogenic fungi. The



IJSURP Publishing Academy

International Journal Of Scientific And University Research Publication
Multi-Subject Journal

Editor.

International Journal Of Scientific And University Research Publication



+965 99549511



+90 5374545296



+961 03236496



+44 (0)203 197 6676

www.ijsurp.com