

International Journal Of Scientific And University Research Publication

ISSN No 301/704

Listed & Index with **ISSN Directory, Paris**



Multi-Subject Journal

Volum : (13) | Issue : 205 |

INTERNATIONAL JOURNAL OF SCIENTIFIC AND UNIVERSITY RESEARCH PUBLICATION



Research Paper



"COMPLEMENTARY MEDICINE": COMPLEMENTARY AND ALTERNATIVE HEALTH APPROACHES IN PEDIATRIC AND ADOLESCENT GYNECOLOGY

Frank M. Biro MD Child Life and Integrative Care		
Division of Adolescent and Transition Medicine		
Cincinnati Children's Hospital Medical Center		
Cincinnati		
Ohio.		
ABSTRACT Complementary and alternative health care approaches are prevalent in the patients and families served by practitioners in pediatric and adolescent gynecology. This article		

addresses gaps in knowledge, including new terminology, prevalence of use, rates of and reasons behind nondisclosure, and potential interactions of herbal products with prescribed medication. It closes with practical complementary health approaches to the adolescent with dysmenorrhea.

KEYWORDS: Complementary medicine, Complementary health approaches, Dysmenorrhea,

INTRODUCTION

Complementary and alternative health care approaches are prevalent in the patients and families served by practitioners in pediatric and adolescent gynecology (PAG). In the past decades there were guidelines proposed by a national committee regarding the recommended competencies in complementary and alternative health approaches, which included delineation of values, knowledge, attitudes, and skills.1 However, a survey of US medical school curricula in 2013 noted that 49% of schools did not have a clerkship or course available, and of those schools who had offered, the most frequent curricular offering was an elective course (71%).2 In this article we address gaps in knowledge, including new terminology, prevalence of use, rates of and reasons behind nondisclosure, and potential interactions of herbal products with prescribed medication.

We close with introducing a second competency, complementary health approaches (CHA) to the adolescent with dysmenorrhea.

Definitions

There have been several changes in terminology regarding complementary and alternative medicine. The National Institutes of Health program that oversees funding opportunities has been renamed the National Center for Complementary and Integrative Health, and the center has recommended modifications in the terms used to describe these practices, as summarized on their Web site (https:// nccih.nih.gov/health/integrative-health). For nonmain-stream practices used together with conventional medicine,

Drs Biro and Bloemer indicate no conflicts of interest.

* Address correspondence to: Frank M. Biro, MD, Child Life and Integrative Care, Division of Adolescent and Transition Medicine, Cincinnati Children's Hospital Medical Center, 3333 Burnet Ave, Cincinnati, OH 45229; Phone: (513) 636-8580, Email address: frank.biro@cchmc.org (F.M. Biro).

the term "complementary" is recommended; this is con- trasted to "alternative" when nonmainstream practices are used in place of conventional medicine. CHA are practices and products of nonmainstream origin, and "integrative health" is defined as complementary approaches incorporated into mainstream health care.

The National Center for Complementary and Integrative Health has identified 2 broad subgroups of CHAdnatural products (herbs, vitamins and minerals, probiotics) and mind and body practices (chiropractic and osteopathic manipulation, yoga, meditation, massage therapy); there are "other" approaches, such as traditional healers, Ayurvedic medicine, traditional Chinese medicine, not included in those subgroups. A group of researchers have proposed 3 categories: CHA products, practices, and providers, 3 terminology we will use in the remainder of this article (Table 1).

Adults

CHA is widely used in the United States. The National Center for Health Statistics has performed the National Health Interview Survey (NHIS), a nationally representative, cross-sectional study conducted through household inter- view surveys. The Adult Alternative Medicine supplement was conducted in 2002, 2007, and 2012, and asked about use of specific CHA over the previous 12 months. The survey defined "adult" as ages 18 years and older, whereas "pediatric" was defined as ages 4-17 years. Because the pediatric and adolescent age range includes pediatric and young adult, as defined in the NHIS, this article examines CHA use among adult and pediatric ages. According to the 2012 NHIS survey, 33.2% of adult Americans used 1 or more CHA.4

Nonvitamin, nonmineral dietary supplements were the commonly used CHA approach, used by 17.7% of American adults. Of these products, fish oil was the single most common supplement (7.8% overall), followed byF.M. Biro, N.L. Bloemer / J Pediatr Adolesc Gynecol 32 (2019) 3e6 nondisclosure to their primary provider, 42% did not traditional healer, massage therapy included the provider had less knowledge of CHA type CHA, complementary health approaches.Adapted from Upchurch and Rainisch.

Table 1 Categories of CHA	disclose, and were more likely not to disclose yoga (65%)
CHA Modality	Examples
Products	Nonvitamin, nonmineral dietary supplements including fish
Practices	oil, melatonin, glucosamine, and probiotics Yoga, tai chi, qi gong, guided imagery, stress management techniques
Providers	Chiropractic/osteopathic manipulation, acupuncture,

glucosamine (2.6%), and probiotics (1.6%). The second most commonly used approach was deep breathing exercises (10.9%) followed by practices of yoga/tai chi/qi gong (10.1%). The peak in any CHA use was reported in the 2007 survey (35.5%).4

<u>Children</u>

The past 2 NHIS surveys asked parents about CHA use in their children 17 years and younger. Parents reported 11.6% used 1 or more approaches in the previous 12 months, with no change in prevalence between the 2007 and 2012 sur- veys. The most common uses in the pediatric ages were nonvitamin, nonmineral dietary supplements (4.9%), chiropractic/osteopathic manipulation (3.3%), and yoga/tai chi/qi gong in 3.2%. Of the dietary supplements, fish oil and melatonin were most commonly used.5

Reasons for Use of CHA

Several investigators have examined reasons for CHA use. Adult CHA users report that these approaches improved overall health (76%), were natural (62%), reduced stress (58%), and they can use it on their own (53%).6 Most users identified wellness or wellness and treatment as the reason they used CHA, whereas 45% reported using CHA for treatment of specific conditions.7 In an earlier NHIS survey, the conditions for which CHA was used most frequently were back pain, head cold, neck or joint pain, arthritis, and anxiety or depression.8 Examining data from the 2007 NHIS reported by parents on use by pediatric patients, adoles- cents were more likely that infants or toddlers to use CHA.9

Adjusting for multiple variables, pediatric/adolescent users were more than 4 times as likely to have a parent with a college education and 1.6 times as likely to use prescription medication. Pediatric users were more likely to have anxi- ety or stress, or dermatologic or musculoskeletal conditions. Use of CHA by a parent increased likelihood of use by their child nearly fourfold.9

Nondisclosure of CHA

In the 2012 NHIS survey, 63% of adults disclosed use of any CHA to their providers, and the rates of disclosure were dependent on the type of CHA.6 That is, 55% disclosed use of CHA providers, 73% disclosed use of CHA products, and 46% disclosed use of CHA practices; disclosure was more likely if the individual experienced improved coping and well- being, or improved health.6 When examining patterns of (7.6%), they were concerned about a negative reaction from the provider (3.0%), they were worried the provider would discourage that use (2.8%), and the provider previously had discourage use of CHA (2.0%).10 Of note, in a survey of patients of children with developmental disabilities, 23% of patients did not disclose CHA use, contrasted to 20% of parents overall.11 Reasons for nondisclosure were that the provider did not ask (61%), followed by parent did not think the provider needed to know (47%).11

HerbeDrug Interactions

Nondisclosure is especially important because of herbedrug interactions and specific organ toxicity that might accompany herbal and dietary supplements. There are multiple mechanisms of herbedrug interactions, as reviewed by Fasinu et al.12 For example, several herbs are, or could have the potential to be, pharmacologic precursors. Herbal and dietary supplements could affect drug absorption, or induce or inhibit metabolic enzymes and transport proteins, or alter renal excretion of drugs and metabolites, leading to either pharmacotoxicity or treatment failure.12 A review of druginduced liver injury reported that herbal and dietary supplements were implicated in 19% of cases, and noted that anabolic steroids were associated with cholestatic hepatic injury, whereas green tea extract was associated with acute hepatitis.13 Of special interest to the PAG provider, the 2 herbal products most commonly used concurrent with prescription medicines are ginkgo and St John's wort, and both can interfere with prescription medicines.14 An evidencebased review of St John's wort and oral contraceptives noted that St John's wort was a potent inducer of cytochrome P450 3A-4 enzyme, and was associated with increased risk of ovulation and breakthrough bleeding, suggesting decreased contraceptive efficacy.15 Overall, studies involving herbedrug interactions have been limited to a small number of conventional and herbal drugs.14 Additionally, studies might also differ ac- cording to the specific formulation used. For example, herbal products vary in composition, and a single product might have varying mixtures and concentrations of the active phytochemicals within that given product.16

Practical Applications

These data would recommend the importance of speaking with patients and their families about CHA, to facilitate safe and effective health care. Incorporating conversations around CHA usage into clinical interactions could initially appear challenging. The provider should focus on 3 issues when framing questions intended to elicit accurate F.M. Biro, N.L. Bloemer / J Pediatr Adolesc Gynecol 32 (2019) 3e6 and important information about patient CHA usage. Acknowledging the following is a broad generalization, CHA are wellness oriented,17 holistic in scope (specifically including body, mind, emotions, and spirit), and intended to foster patient empowerment. Examples of wellness ori- ented questions include: "When do you feel your best?" "What types of products, practices, or practitioners do you use for your overall wellness or to help with your symp- toms?" Use of open-ended questions will invite shared information regarding physical, mental, emotional, and spiritual approaches the patient and family find helpful, and might help to elicit a more holistic picture of the patient's wellness perspectives and practices.

Because overall "wellness" is a subjective experience, many CHA intentionally emphasize the patient and their family as the authorities (the "author" of the experience) on what exactly wellness is, and how to attain and maintain it. An approach that implies "teach me what works (and what doesn't) for you" affirms for the patient the importance of their subjective experience, and clarifies the willingness of the health care provider to hear it, while fostering a sense of empowerment. For example, when the PAG provider explains their concerns about specific drugesupplement interactions; or explains why questions about a particular style of yoga might be related to a patient's pain issues; or why it might be advisable to seek out a mental health professional before going on a month-long meditation retreat for treatment of depression. "Integrative medicine" implies an integration of the expertise of every one of the parties involved, whether conventional or complementary health care products, practices, and providers, or the subjective perspectives of the patient and their family. With overall wellness, holistic orientation and patient empowerment as guiding principles, eliciting comprehensive and imperative details about CHA usage allows such "integration" toward safe and effective health care.

CHA for Dysmenorrhea

Dysmenorrhea is one of the most common concerns in adolescent women, with a prevalence of 16%-93%.18

Although effective modalities exist (specifically, nonste- roidal antiinflammatory drugs [NSAIDs] and combined oral contraceptives), up to one-fourth of adolescents might not respond,18 leading many parents and patients to inquire about "natural" approaches either before implementation, or as augmentation, to NSAIDs or oral contraceptives. The literature reports several potentially effective CHA modalities for dysmenorrhea, including local heat, transcutaneous nerve stimulation, acupuncture, behavioral interventions, and dietary and herbal supplements.

A Cochrane Collaborative Review from 2002 noted that transcutaneous nerve stimulation performed better than placebo, with one-third of patients responding; of note, few studies included adolescents.19

A meta-analysis of acupuncture for dysmenorrhea examined 60

IF: 4.176 | IC Value: 78.46

randomized control trials, published before December 2017. Most of the studies examined women in their late teens to twenties, and concluded that acupuncture reduced menstrual pain and associated symptoms more effectively than no treatment or NSAIDs.20 The authors explained that acupuncture in traditional health approaches helped the body circulate blockages in energy, termed qi. An allopathic explanation for acupuncture is the stimulation of the nervous system by antidromic axon reflexes, with the release of serotonin and exogenous opioids. The authors recommend more rigorously designed trials to address quality and methodology.20

A Cochrane collaborative review from 2011 examined behavioral interventions for dysmenorrhea; most studies involved participants who were college-aged or older.21 The review noted that several different behavioral interventions have been used for dysmenorrhea, which include modifying behavior, or thoughts or cognitions, and physical and cognition procedures such as biofeedback, desensitization, relaxation training, and hypnotherapy. The investigators ultimately identified 13 trials, of which 5 met study criteria. The authors concluded that relaxation techniques with or without imagery, pain management training, plus relaxation with biofeedback might help with dysmenorrhea. They recommend future randomized controlled trials with adequate sample size, use of objective measurement scales, and comparative studies against standardized treatments (such as NSAIDs) to help address the utility of these approaches.21

There was an updated Cochrane collaborative review on dietary supplements,22 which excluded several studies incorporated into the earlier review.23 The more recent review notes there was limited evidence of benefit for several dietary supplements, including fenugreek, ginger, valerian, zetaria, zinc sulfate, fish oil, and vitamin B1. Although most of the studies were implemented in women 18-30 years of age, several included adolescents. The authors believed that there was no high-quality evidence because of small sample sizes and inconsistencies; they did note more research was justified. Of note, a recent international publication examined the combination of 2 interventions previously believed to have benefit for management of dysmenorrhea: fish oil24 and vitamin B1 (thiamine).25

Students received placebo, 100 mg vitamin B1 alone, 500 mg fish oil alone, or vitamin B1 and fish oil. The combination of B1 and fish oil was most effective; and B1 alone was next most effective and exhibited the fewest side effects. Fish oil alone was more effective than placebo, but both interventions with fish oil had greater rates of side effects.26

CONCLUSION

We reviewed the new terminology regarding CHA, prevalence of use, rates of and reasons for nondisclosure, potential interactions of complementary approaches with prescribed medication, and potentially effective CHA to the adolescent with dysmenorrhea. CHA are quite common, with a minimum prevalence of one-third of patients.

However, nondisclosure rates could be 40% or greater. One of the major issues of nondisclosure is the potential interaction with prescribed medication, or direct toxicity of the agent. We encourage PAG professionals to provide an openended, nonjudgemental style when eliciting information about complementary and integrative health approaches in the PAG patient and their family.

ref_str

- 1. Kligler B. Maizes V. Schachter S. et al: Core competencies in integrative medicine for medical school curricula: a prop Acad Med 2004; 79:521
- 2. Cowen VS. Cyr V: Complementary and alternative medicine in US medical schools. Adv Med Edu Prac 2015; 6:113
- 3. Upchurch DM, Rainisch BK: A sociobehavioral model of complementary and alternative medicine providers, products, and practices: findings from the National Health Interview Survey, 2007, Evid Based Complement Alternat Med 2012: 18:100
- 4. Clarke TC, Black LI, Stussman BJ, et al: Trends in the use of complementary health approaches among adults: United States, 2002-2012, National health statistics reports: no 79. Hyattsville, MD, National Center for Health Statistics, 2015
- 5. Black LI, Clarke TC, Barnes PM, et al: Use of complementary health approaches among children aged 4-17 years in the United States: National Health Interview Survey, 2007-2012. National health statistics reports; no 78. Hyattsville, MD, National Center for Health Statistics, 2015
- 6. Sirois FM, Riess H, Upchurch DM: Implicit reasons for disclosure of the use of complementary health approaches (CHA): a consumer commitment perspective. Ann Behav Med 2017; 51:764
- 7. Stussman BJ, Black LI, Barnes PM, et al: Wellness-related use of common complementary health approaches among adults: United States, 2012. National health statistics reports; no Hyattsville, MD, National Center for Health Statistics, 2015
- 8. Barnes PM, Powell-Griner E, McFann K, et al: Complementary and alternative medicine use among adults: United States, 2002. Advanced Data from Vital and Health Statistics; no 343. Hyattsville, MD, National Center for Health Statistics, 2004
- 9. Birdee GS, Phillips RS, Davis RB, et al: Factors associated with pediatric use of complementary and alternative medicine. Pediatrics 2010: 125:249
- 10. Jou J. Johnson PJ: Nondisclosure of complementary and alternative medicine use to primary care physicians: findings from the 2012 National Health Interview Survey. JAMA Intern Med 2016: 176:545
- 11. Lindly O, Thorburn S, Zuckerman K: Use and nondisclosure of complementary health approaches among US children with developmental J Dev Behav Pediatr 2018; 39:217
- 12. Fasinu PS, Bouic PJ, Rosenkranz B: An overview of the evidence and mechanisms of herbedrug interactions. Front Pharmacol 2012; 69:1
- 13. Navarro VJ, Khan I, Bjo€rnsson E, et al: Liver injury from herbal and dietary supplements. Hepatology 2017; 65:363
- 14. Choi JG, Eom SM, Kim J, et al: A comprehensive review of recent studies on herb-drug interaction: a focus on pharmacodynamic intera J Altern Complement Med 2016; 22:262
- 15. Berry-Bibee EN, Kim MJ, Tepper NK, et al: Co-administration of St. John's wort and hormonal contraceptives: a systematic review. Contraception 2016; 94:668
- 16. Mok DK, Chau F: Chemical information of Chinese medicines: a challenge to Chemometr Intell Lab Syst 2006; 82:210
- 17. Upchurch DM. Rainisch BW: The importance of wellness among users of complementary and alternative medicine: findings from the 2007 National Health Interview Survey. BMC Complement Altern Med 2015: 15:362
- 18. DeSanctis V, Soliman A, Bernasconi S, et al: Primary

IF: 4.176 | IC Value: 78.46

dysmenorrhea in adolescents; prevalence, impact and recent knowl Ped Endocrinol Rev 2015; 13:465

- <u>19. Proctor M, Farquhar C, Stones W, et al: Transcutaneous electrical</u> nerve stimulation for primary dysme Cochrane Database Syst Rev 2002;1:CD002123
- 20. Woo HL, Ji HR, Pak YK, et al: The efficacy and safety of acupuncture in women with primary dysmenorrhea. A systematic review and meta-analysis. Medicine (Baltimore) 2018; 97:e11007
- 21. Proctor M, Murphy PA, Pattison HM, et al: Behavioural interventions for dysmenorr Cochrane Database Syst Rev 2007; 3:CD002248
- 22. Pattanittum P, Kunyanone N, Brown J, et al: Dietary supplements for dysmenorrhoea. Cochrane Database Syst Rev 2016; 3:CD002124
- 23. **Proctor ML, Murphy PA:** Herbal and dietary therapies for primary and secondary dysmenorrhea. Cochrane Database Syst Rev 2001: 3:CD002124
- 24. Harel Z, Biro FM, Kottenhahn RK, et al: Supplementation with omega-3 poly- unsaturated fatty acids in the management of dysmenorrhea in adolescents, Am J Obstet Gynecol 1996; 174:1335
- 25. Gokhale LB: Curative treatment of primary (spasmodic) dysmenorrhoea. Indian J Med Res 1996; 103:227
- 26. Hosseinlou A, Alinejad V, Alinejad M, et al: The effects of fish oil capsules and vitamin B1 tablets on duration and severity of dysmenorrhea in students of high school in Urmia-Iran. Glob J Health Sci 2014; 6:124



IJSURP Publishing Academy

International Journal Of Scientific And University Research Publication Multi-Subject Journal

Editor.

International Journal Of Scientific And University Research Publication





C +90 5374545296





www.ijsurp.com