



International Journal Of Scientific And University Research Publication

ISSN No **301/704**

Listed & Index with
ISSN Directory, Paris



Multi-Subject Journal



BIOACCUMULATION OF HEAVY METALS IN CHANNA PUNCTATUS (BLOCH) IN RIVER RAMGANGA (U.P.), INDIA

Robeena Sarah || Toxicology Laboratory
Department of Zoology
Govt. Raza P.G. College
Rampur-244901 (U.P.)
India.

Ganga is the largest riverine system of India with a fragile ecosystem. Its prone to anthropogenic disturbances because of its cultural, economic and environmental values. The

contamination of river Ganga by heavy metals (HM) is due to biotic (anthropogenic sources) and abiotic (pesticides, fertilizers) sources that poses a devastating health hazard to human, plant and edible fish life. The chemical analysis with the help of atomic absorption spectrometer performed on its water samples demonstrated the accumulation of heavy metals such as Arsenic (As), Lead (Pb), Cadmium (Cd), Iron (Fe), Zinc (Zn). Moreover, the spectrophotometric analysis indicated clearly the accumulation of heavy metals in order of occurrence (Fe> As> Cd> Zn> Pb) in liver and (Zn> Fe> As> Cd> Pb) in kidney of edible fish *Channa punctatus*. The present study has been used to sensitively monitor the extent of heavy metals pollution in the biotic aqua life of river Ramganga system and its suggested that the bioaccumulation of heavy metal in *Channa punctatus* has reached above permissible limits for human consumption, indicating potential health risks. Necessary biological steps should be taken to handle such food pollution and prevent the environmental risk and food chain disruption.

Bioaccumulation of heavy metals in *Channa punctatus*; Ramganga River,

مقدمة

bodies (Gupta et al. 2016; Ahmad et al. 2019) as well as that of its ecosystem (Dudani et al. 2017; Sharma 2019). Heavy metals thus discharged from the source, leach into underground waters, deposits in the aquifer, or are washed away by run-off into surface waters. Industrial wastes containing heavy metals on entering aquatic environment causes biochemical disturbances in the fish (Ansah et al. 2019; Varol and Sünbül 2019).

The concentrations of heavy metals in aquatic environment and its organisms have been of considerable interest to understand because of their toxic effects (Wani et al. 2017; Satapathy and Panda 2018). Over the last few decades, there has been growing interest in determining heavy metal concentration in the aquatic environment and its measurement of contamination levels in public food supplies, especially fishes (Velusamy et al. 2014; Fatima et al. 2015; Arulkumar et al., 2017; Pal and Maiti 2018; Kumari et al. 2018).

The aim of this study was to evaluate the concentrations of heavy metals i.e. Fe, Zn, Cd and Pb in surface water of river Ramganga as well as groundwater of nearby regions and its influence on the tissues. Further, *Channa punctatus*, of inhabiting fresh water teleost fish, their hazardous levels were compared with available certified safety guidelines proposed by World Health Organization (WHO) and Food C. and Agricultural Organization (FAO) for human consumption was selected in this experiment for bio-monitoring the *punctatus* concentration of heavy metals in this region because of its hardy nature, high nutritional value and its round the year availability.

aterial and MethodsM Description of Study Area

The study area was River Ramganga located in district Moradabad, Uttar Pradesh, India (Fig.1A-B). The study was conducting at five different locations of the river and the nearby regions as described clearly in Figure 1. The selection was based on the severe anthropogenic activities like brass industries which is continuously discharging their effluents into the river and waste scrapping and illegal cremation of the dead bodies by the local people of Katghar, Jama Masjid and its nearby regions. Such activities have increased the concentration of heavy metals in the surface water as well as groundwater. These areas also received the domestic waste water of adjacent communities living therein.

Collection of Samples

Water samples (five per each location) were collected in the month

Water is essential for all forms of life. Clean, clear & odorless freshwater used for drinking and other domestic purposes is the gift from the nature. From the entire water reserve of the earth, approximately 97% of water consists of seas and oceans while as freshwater resources contribute only 3% of water (Shrestha et al. 2017). from the earth's freshwater which is locked up in glaciers and ice caps on poles, 30.1% is groundwater (Adhikari and Mal, 2019). In India, the major source of water used for drinking purposes and domestic needs is the Groundwater. Life cannot be sustained more than few days without water, while an inadequate supply of water may change the pattern of distribution of organisms as well as of human beings. But due to anthropogenic activities, the water necessary for our survival is becoming hazardous every day (Rao et al. 2018).

Ganga is the well-developed largest riverine system of India and the fifth longest in the world (Mukherjee and Pal 2018; Sahu et al. 2018). It has several important cultural, economical and environmental values and provides water for approximately to 450 million people (Behera et al. 2011). Moreover, Ganga River is being used for fishing and aquaculture (Gupta et al. 2018). Fishes in addition to its important source of protein have rich contents of essential minerals, vitamins and unsaturated fatty acids (Medeiros, 2012). They can normally accumulate heavy metals from food, water and sediments as they are situated at the top of aquatic food chain (Afshan et al. 2014). Due to rapid industrialization, this river is being continuously degrading due to direct discharge of industrial wastes, agricultural runoff and anthropogenic activity along the river bank (Jani et al. 2018). These wastes contain health hazard chemicals like salts of chromium, copper, cadmium, arsenic, mercury and lead which interact with aquatic environment and affect the river ecosystem (Usmani and Kumar 2017; Singh et al. 2017, 2018). Heavy metals containing pollutants besides poses serious water pollution problem due to their toxicity, persistence and bioaccumulation, also accumulate in tissues of plants and animals leading to toxicity in aquatic biota (Chopra et al. 2011; Pandey et al. 2014). Fishes and humans being the top consumers in aquatic ecosystem and food chain are badly affected through the accumulation of these heavy metals (Afshan et al. 2014). Heavy metals are the metallic elements which are dense and heavy and are present in trace amounts (Singh et al. 2013; 2018). Heavy metals lead to alteration in physical, chemical & biological properties of water

unnecessary high level of these toxic trace metals which were transferred to man through its consumption (Adeniyi and Yusuf, 2007; Sang et al. 2018; Pal et al. 2018). The content of these toxic heavy metals can counteract their beneficial effects and can lead to several adverse effects to human health (Castro-Gonzalez and Mendez-Armenta, 2008). A known biological function indicates that Cadmium does not break down in the environment, remain in fish body for long periods of time and can bioaccumulate for many years after its exposure to low levels (Wang et al. 2018; Markowicz et al. 2019). Kwaansa-Ansah et al. (2019), proved that the fish liver is the target organ for Fe. Respiratory disruption due to physical clogging of the gills is suggested as a possible mechanism for iron toxicity (Wani et. al, 2017).

Aquatic organisms have tendency to accumulate lead from water and diet, although there is evidence that origin of lead accumulation in fish, is most probably from contaminated water rather than that of diet (Creti et al., 2010; Velusamy et al. 2014; Dudani et al. 2017; Ahmed et al.2019).

The aquatic ecosystem is contaminated by arsenicals and their effect on the aquatic organisms has now emerged as a serious threat to environment (Chopra et al. 2011; Gupta et al. 2016; Dudani et al. 2017). Aquatic ecosystem is too complex and hence indicators are useful for its effective bio-monitoring. The environmental monitoring programmes were evolved to measure impact of stress inducers on aquatic fauna (Kumar et al. 2010; Chopra et al. 2011). Fish is a useful bio-indicator for the determination of heavy metal pollution in aquatic ecosystems Sayegh Petkovsek et al. 2011). Pal et al. (2014) and Punetha et al. (2015) found that Moradabad district also had high concentrations of Cd in its water samples because these areas are densely populated, industrialized and are hubs of electronic industries, so the possibility of Cadmium accumulation is high in this region.

استنتاج

The purpose of this investigation is to detect the concentration of heavy metals i.e. Fe, Cd, As, Pb and Zn in river Ramganga of district Moradabad, Uttar Pradesh (India). Results showed that the river is highly contaminated with heavy metals accumulated in water as well as fish. The values of Fe, As, Cd and Pb detected was more than the maximum permissible limit when compared with the National and International Organizations like B.I.S and W.H.O respectively while that of Zn is under the permissible limit. As we know that Moradabad is a famous city for its brass work. The cause of such accumulation is mainly due to illegal dumping of wastes from various small-scale industries, scrapping and illegal burning of wastes near the banks of the river.

Hence, water is not suitable for fish and human consumption. On the other hand, Fe, Zn and Pb were found to be under the Maximum Permissible Limit (MPL) of W.H.O. (1985) and FEPA (1991) in fish tissue samples while that of Cd and As is above the MPL range. Therefore, results suggested that Cadmium and Arsenic is accumulating in the tissues of fish through contaminated river water and thus, it might be transferring via food chain to humans. So it is necessary to monitor the quality of water at regular intervals and more

of September and October (2017) from the river and the nearby regions. One liter sterilized bottles were used to collect water samples for heavy metal analysis. The six-medium sized fresh water fish,) samples (five for each location) were collected *punctatusChanna* (with the help of local fisherman from the affected regions.

ample preparation for Heavy metal analysisS

Water samples collected were filtered and acidified with Nitric acid and stored for heavy metal analysis. These six medium sized fish samples were sacrificed and the main target organ was removed. Fish tissue samples were dried at 1050C and powdered using porcelain pestle and a mortar. Weighted amount of dried tissues was digested with a mixture of nitric acid and perchloric acid in the ratio of 3:1. Digested samples were then filtered with whatman filter paper and made up to 25.0 ml with metal free double distilled water (Tabassum et al. 2016). The metals in fish and water samples were analyzed using Atomic Absorption Spectrometer and results are expressed in µg/g (for tissue samples) and mg/L (for water samples).

atistical AnalysisSt

Statistical analysis of the five-sampled data was done analysis variance (ANOVA) using a statistical package, SPSS 22.0 (IBM Corporation, Chicago, IL, USA). The significance of the results was ascertained at P<0.05.

Results

The heavy metals detected in water were in [Heavy metals in water:](#) the decreasing order Fe> Zn> Pb> As> Cd. The values of Fe, As, Cd and Pb detected was more than the maximum permissible limit when compared with the National and International Organizations like B.I.S and W.H.O respectively while that of Zn was under the permissible limit. Hence, water is not suitable for fish and human consumption. Results shown in Table 1 depicts the range, mean and standard deviation of heavy metal content in water whereas Table 2 shows the comparison of obtained values of water samples with W.H.O. and B.I.S. values.

ioaccumulation Factor (BAF)B

Bioaccumulation Factor (BAF) is calculated as the ratio of concentration of a pollutant (heavy metal) accumulated in the tissue of an organism with respect to the concentration of that pollutant i.e. heavy metal in water body (Authman and Abbas, 2007). The order of bioaccumulation in fish liver was Fe>As>Cd>Zn>Pb while that in kidney Zn>Fe>As>Cd>Pb. The values observed during the present study when compared with the recommended values for heavy metals i.e. Fe, Zn and Pb were found to be under the Maximum Permissible Limit (MPL) of W.H.O. (1985) and FEPA (1991) while that of Cadmium is found to be 0.2mg/kg and Arsenic is above 0.26mg/kg recommended maximum permissible limit. Hence, the results suggested that Arsenic and Cadmium has been accumulating in the tissues of fish inhabiting river Ramganga.

Discussion

Presence of heavy metals in the environment can cause serious problems to all organisms and its bioaccumulation in the food chain can be highly dangerous to human health (Devi and Yadav 2018; Adimalla and Wang 2018; Ahmed et al., 2019). It can also possible that environmental toxicants may increase the susceptibility of aquatic animals to various diseases by interfering with the normal functioning of their immune, reproductive and developmental processes (Rezvanfar et al. 2016; Sang et al. 2018). Concentration of heavy metals in different tissues/organs of fishes is directly influenced by contamination in aquatic environment, its up- take, regulation and elimination inside the fish body (Yang et al. 2016; Brraich et al. 2017). Fish have ability to concentrate metals in their muscles (Zhaoa et al. 2018; Juncos et al 2019) and as they are a part of human nutrition. They need to be carefully screened to ensure

- Physicochemical **Gupta A., Bhatnagar P. and Bakre P. P. (2016).**..13
properties of water and heavy metals (Lead and Zinc) in water and
sediment of a reservoir and drainage of Jaipur (Rajasthan) – A
comparative study. International Journal of Fisheries and Aquatic
Studies 2016; 4(5):
407-412.
<http://www.fisheriesjournal.com/archives/2016/vol4issue5/PartF/4-5-979.pdf>.6.
- Taxonomic **Gupta D., Dwivedi A.K. and Tripathi M. (2018).**..14
validation of five fish species of subfamily Barbinae from the Ganga
river system of northern India using traditional and truss analyses.
PLoS ONE 13(10):
<https://doi.org/10.1371/journal.pone.0206031>.e0206031.
- Jani K., Ghattargi V., Pawar S., Inamdar M., Shouche Y. and**..15
Anthropogenic activities induce depletion in **Sharma A. (2018).**
microbial communities at Urban sites of the River Ganges. Curr
<https://doi.org/10.1007/s00284-017-1352-5>. Microbiol 75: 79-83.
- Juncos R., Arcagni M., Squadrone S., Rizzola A., Arribérea M.,**..16
Campbell L.M., Brizioc P., Abetec Barriga J.P., Battin M.A.,
Interspecific differences in the **M.C., Guevara S. R. (2019).**
bioaccumulation of arsenic of three Patagonian top predator fish:
Organ distribution and arsenic speciation. Ecotoxicology and
Environmental Safety, 168: 431-442.
<https://doi.org/10.1016/j.ecoenv.2018.10.077>
- Assessment of **Kumari P., Chowdhury A. and Maiti S. K. (2018).**..17
heavy metal in the water, sediment, and two edible fish species of
Jamshedpur Urban Agglomeration, India with special emphasis on
human health risk, Human and Ecological Risk Assessment: An
International Journal, 24(6): 1477-1500. DOI:
10.1080/10807039.2017.1415131.
- Kwaansa-Ansah E. E., Nkrumah D., Nti S. O. and Opoku F. (2019).**..18
Adsorption of Heavy Metals (Cu, Mn, Fe and Ni) from
Scales. Pollution, *Oreochromis niloticus* surface water using
5(1):115-122. DOI: 10.22059/poll.2018.259347.454.
- Markowicz F., Król G. and Szymańska-Pulikowska A. (2019).**..19
Biodegradable package- Innovative purpose or source of the problem.
Journal of Ecological Engineering, 20(1):
228-237.<https://doi.org/10.12911/22998993/94585>.
- Channel migration zone mapping **Mukherjee K. and Pal S. (2018).**..20
of the River Ganga in the Diara surrounding region of Eastern India.
Environ Dev Sustain, 20:
<https://doi.org/10.1007/s10668-017-9984-y>.2181-2203.
- Seasonal variation of heavy metals in **Pal D. and Maiti S. K. (2018).**..21
Labeo rohita water, sediment, and highly consumed cultured fish (
) and potential health risk assessment in aquaculture *Labeo bata* and
pond of the coal city, Dhanbad (India). Environmental Science and
Pollution Research 25:
<https://doi.org/10.1007/s11356-018-1424-5>.12464-12480.
- Assessment of heavy **Pal R., Gupta M. A. and Tripathi A. (2014).**..22
metals in suspended particulate matter in Moradabad, India. Journal
of Environmental Biology, 35: 357-361.
- Pereira R., Leite E., Raimundo J., Guilherme S., Puga S., Ribeiro**..23
F. P., Santos M. A., Canário J., Almeida A. , Pacheco M. and
Metals (loids) targeting fish eyes and brain in a **Pereira P. (2018).**
contaminated estuary-Uncovering neurosensory (un)susceptibility
through bioaccumulation, antioxidant and morphometric profiles.
Marine Environmental Research,140: 403-411.
- De Silva B. C.J., Hossain S., Dahanayake P. S. and Heo G. J. (2018).**..24
Frozen white-leg shrimp (*Litopenaeus vannamei*) in Korean
markets as a source of *Aeromonas* spp. harboring antibiotic and heavy
metal resistance genes. Microbial Drug Resistance, 24(10):1587-
1598. DOI: 10.1089/mdr.2018.0035.
- Application of multivariate statistics **Adhikari K. and Mal U. (2019).**..25
in the analysis of groundwater geochemistry in and around the open
cast coal mines of Barjora block, Bankura district, West Bengal,
India. Environmental Earth Sciences, (2019)
<https://doi.org/10.1007/s12665-019-8071-0>.78:72.
- Punetha D., Tewari G., Pande C., Kharkwal G. C. and Tewari K. (2015).**..26
Investigation on heavy metal content in common grown
vegetables from polluted sites of Moradabad district, India. Journal of
the Indian Chemical Society, 92(1): 97-103.
- Rao N. S., Sunitha B., Rambabu R., Rao P. V. N., Rao P. S., B. D. (2018).**..27
Quality and **Spandana, Sravanthi M. and Marghade D. (2018).**
degree of pollution of groundwater, using PIG from a rural part of
Telangana State, India. Applied Water Science,
[018-0864-x](https://doi.org/10.1007/s13201-018-0864-x).<https://doi.org/10.1007/s13201-018-0864-x>.8:227
- research work is needed to find suitable ecofriendly and
cheaper antagonist of heavy metal pollution.
- The authors declare that they **Conflict of Interest:**
“All **Ethical approval:** have no conflict of interest.
applicable international, national, and/or institutional
guidelines for the care and use of animals were followed by
the authors.”
- The authors would like to extend their **Acknowledgment**
sincere appreciation to the Deanship of Scientific Research at
king Saud University for its funding this Research group NO
(RG-1435-014).
- ref_str
- Distribution, contamination, and **Adimalla N. and Wang H. (2018).**..1
health risk assessment of heavy metals in surface soils from northern
Telangana, India. Arabian Journal of Geosciences,11: 684.
<https://doi.org/10.1007/s12517-018-4028-y>
- Assessment of **Ahmad R., Kaushik H. and Ranjan R. K. (2019).**..2
microbial communities and heavy metals in urban soils of Patna,
Bihar (India). Arabian Journal of Geosciences, 12:
<https://doi.org/10.1007/s12517-018-4188-920>.
- Ahmed S., Khurshid S., Qureshi F., Hussain A. Bhattacharya A. (2019).**..3
Heavy metals and geoaccumulation index development for (2019),
groundwater of Mathura city, Uttar Pradesh. Desalination and
Water Treatment Journal, 138: 291-300. DOI:
10.5004/dwt.2019.23322
- Ansah K. E. E., Nkrumah D., Nti S. O. and Opoku F. (2019).**..4
Adsorption of heavy metals (Cu, Mn, Fe and Ni) from surface water
using *Oreochromis niloticus* Scales. Pollution, 5(1):
115-122.DOI:10.22059/poll.2018.259347.454
- Toxic **Arulkumar A., Paramasivam S. and Rajaram R. (2017).**..5
heavy metals in commercially important food fishes collected from
Palk Bay, Southeastern India. Marine Pollution Bulletin,119: 454-459.
<https://doi.org/10.1016/j.marpolbul.2017.03.045>
- Accumulation and **Authman M. M. N. and Abbas H. H.H. (2007).**..6
distribution of copper and zinc in both water and some vital tissues of
) of Lake Qarun, *Mugil cephalus* and *Tilapia zillii* two fish species (
Fayoum Province, Egypt. Pakistan Journal of Biological Sciences, 10:
2106-DOI: 10.3923/pjbs.2007.2106.2122
- Histopathological alterations in **Brraich O. S. and Kaur, M. (2017).**..7
the gills of *Labeo rohita* (Hamilton-Buchanan) due to Lead toxicity.
Indian Journal of Experimental Biology,
<http://nopr.niscair.res.in/handle/123456789/42552>.55(8):576-583.
- Chopra A. K., Sharma M.K. and Chamoli S. (2011).**..8
Bioaccumulation of organochlorine pesticides in aquatic system-an
overview Environ Monit Assess, 173: 905-916. DOI:
10.1007/s10661-010-1433-4
- Chemometric evaluation of heavy **Devi N.L. and Yadav I.C. (2018).**..9
metal pollutants in Patna region of the Ganges alluvial plain, India:
implication for source apportionment and health risk assessment.
Environ Geochem Health, 40(6): 2343-2358. doi:
10.1007/s10653-018-0101-4.
- Dudani S. N., Lakshmapurkar J., Gavali D. and Patel T. (2017).**..10
Heavy metal accumulation in the mangrove ecosystem of south
Gujarat coast, India. Turkish Journal of Fisheries and Aquatic
Sciences 17: 755-766. DOI: 10.4194/1303-2712-v17_4_11.
- Fatima M., Usmani N., Firdaus F., Zafeer M. F., Ahmad S.,**..11
Akhtar K., Husain S.M. D., Ahmad M. H., Anis E. and Hossain
In vivo induction of antioxidant response and **M. M. (2015).**
oxidative stress associated with genotoxicity and histopathological
alteration in two commercial fish species due to heavy metals exposure
in northern India (Kali) river. Comparative Biochemistry and
Physiology, (176–177): 17-30.
<https://doi.org/10.1016/j.cbpc.2015.07.004>.
- Federal Environmental Protection Agency (FEPA) Act (1991).**..12
Guidelines and standards for Industrial effluent, gaseous emissions
and hazardous waste management in Nigeria. National Environmental
Protection Regulations, Federal Republic of Nigeria. Supplement to
Official Gazette Extraordinary - Part B. 78 (42): B15 –31.

- .Growing **Rezvanfar M. A., Hodjat M., Abdollahi M. (2016)**. .28
knowledge of using embryonic stem cells as a novel tool in
developmental risk assessment of environmental toxicants. Life
<https://doi.org/10.1016/j.lfs.2016.05.027>. Sciences, 158: 137-160.
- Sone megafan: a non-**Sahu S., Saha D. and Shukla R. R. (2018)**. .29
imalayan megafan of craton origin, forming a potential groundwater
reservoir in marginal parts of Ganga Basin, India. Hydrogeology
<https://doi.org/10.1007/s10040-018-1829-8>. Journal, 26: 2891-2917.
- Developmental **Sang W., Xu J., Bashir M. H. and Ali S. (2018)**. .30
to heavy metals transferred *Cryptolaemus montrouzieri* responses of
across multitrophic food chain. Chemosphere,
205:690-697. <https://doi.org/10.1016/j.chemosphere.2018.02.073>.
- Source identification, **Satapathy S. and Panda C.R. (2018)**. .31
environmental risk assessment and human health risks associated with
toxic elements present in a coastal industrial environment, India.
Environ Geochem Health, 40(6): 2243-2257. doi:
10.1007/s10653-018-0095-y.
- Risk assessment and mitigation Measures on the **Sharma D. S. (2019)**. .32
heavy Metal polluted water and sediment of the Kolleru Lake in
Andhra Pradesh, India. Pollution, 5(1): 161-178. DOI:
10.22059/poll.2018.263546.493.
- Assessing climate change **.K., Du X. and Wang J. (2017). Shrestha N.** .33
impacts on fresh water resources of the Athabasca River Basin,
Canada. Science of the Total Environment, 601-602:425-440.
- Assessment **Singh H., Kushwaha A. and Shukla D. N. (2018)**. .34
of ecoenvironmental geochemistry of heavy metals pollution of the
river Gandak, a major tributary of the river Ganga in Northern India.
AIP Conference Proceedings 1952, 020038
<https://doi.org/10.1063/1.5032000>. (2018);
- Singh H., Pandey R., Singh S.K. and Shukla D.N. (2017)**. .35
Assessment of heavy metal contamination in the sediment of the
River Ghaghara, a major tributary of the River Ganga in Northern
India. Appl Water Sci, 7: 4133-4149. doi
10.1007/s13201-017-0572-y.
- Singh, A. N., Shrivastava R., Mohan D. and Kumar, P. (2018)**. .36
Assessment of spatial and temporal variations in water quality
dynamics of River Ganga in Varanasi. Pollution, 4(2): 239-250. DOI:
10.22059/poll.2017.240626.310.
- Metal bioaccumulation in tissues of **Usmani Z. and Kumar V. (2017)**. .37
Puntius sarana and Labeo rohita and its associated risk status: A case
study of Damodar River, India. Desalination and Water Treatment,
76: 196-211. doi:10.5004/dwt.2017.20719.
- Environmental contaminants in **Varol M. and Sünbül M. R. (2019)**. .38
fish species from a large dam reservoir and their potential risks to
human health. Ecotoxicology and Environmental Safety, 169:
<https://doi.org/10.1016/j.ecoenv.2018.11.060>. 507-515.
- Velusamy A., Kumar P. S., Ram A. and Chinnadurai S. (2014)**. .39
Bioaccumulation of heavy metals in commercially important marine
fishes from Mumbai Harbor, India Marine Pollution Bulletin, 81:
<https://doi.org/10.1016/j.marpolbul.2014.01.049>. 218-224.
- Breakdown of Fermi **Wang J. R., Liu G. Z. and Zhang C. J. (2018)**. .40
liquid theory in topological multi-Weyl semimetals. Phys.Rev.B,
<https://doi.org/10.1103/PhysRevB.98.205113>. 98:205113.
- Evaluation of **Wani, MA, Shammi, Q. J. and Singh, J. (2017)**. .41
toxicity of heavy metal iron on a fresh water fish Labeorohita and its
behavioural impacts. Research Journal of Pharmaceutical, Biological
and Chemical Sciences, 8(5): 197-203
- Guidelines for Drinking Water Quality. Vol. 5, World **WHO. (1985)**. .42
Health Organization, Geneva, Switzerland.
- Cadmium potentiates **Yang Y., Ye X., He B. and Liu J. (2016)**. .43
toxicity of cypermethrin in zebrafish. Environ Toxicol Chem., 35(2):
435-45. doi: 10.1002/etc.3200.
- Zhaoh H., Quana W., Bekelea T. G., Chena M., Zhanga X. and** .44
Effect of copper on the accumulation and elimination **Qu B. (2018)**.
kinetics of fluoroquinolones in the zebrafish (Danio rerio).
Ecotoxicology and Environmental Safety 156:
<https://doi.org/10.1016/j.ecoenv.2018.03.025>. 135-140.



IJSURP Publishing Academy

International Journal Of Scientific And University Research Publication
Multi-Subject Journal

Editor.

International Journal Of Scientific And University Research Publication



+965 99549511



+90 5374545296



+961 03236496



+44 (0)203 197 6676

www.ijsurp.com