



International Journal Of Scientific And University Research Publication

ISSN No **2364\2018**

Listed & Index with
ISSN Directory, Paris



Multi-Subject Journal



A STUDY ON THE ACUTE TOXICITY OF TRIAZOPHOS TO BRANCHIURA SOWERBYI (BEDDARD, 1982) AND THEIR BEHAVIOURAL CHANGES

Scientist Mariane

ABSTRACT

ABSTRACT The study was made to evaluate the acute toxic effects of Triazophos to benthic oligochaete worm, *Branchiura sowerbyi* along with the changes in their behaviour

during the experiment. The 24, 48, 72 and 96 h LC50 values of Triazophos to *Branchiura sowerbyi* were 8.17, 7.07, 5.81 and 5.04 mg/l respectively. The mortality rate of *Branchiura sowerbyi* varied significantly ($p < 0.05$) with the increasing concentrations of the toxicant from 5.92, 5.60, 4.80 and 4.32 mg/l at 24, 48, 72 and 96h of exposure respectively. A significant variation ($p < 0.05$) on the mortality rate was also recorded at each concentration (from 5.60 mg/l and above) with the increasing exposure time (24, 48, 72 and 96h). The clumping tendency and movement of *B. sowerbyi* were decreased with the increasing concentration of the test chemical and exposure time. The mucous secretion of the worm was increased with the increasing concentration and the time of exposure.

KEYWORDS :

INTRODUCTION

The pesticides are used in agricultural field to eradicate different pest populations for increasing food production (Naveed et al., 2011). When an insecticide is used in the field, it affects its target pests only by 0.1%, but the rest of the insecticide spreads and contaminates the natural ecosystem (Hart and Pimentel, 2002; Mahboob et al., 2011). Triazophos, a newly introduced non-systemic, broad spectrum organophosphate insecticide is used to control different types of pest populations in the field of rice and cotton of all parts of the world (Worthing and Hanrae, 1991; Mingzhang et al., 2003). It is neurotoxic in nature and leads to accumulate the neurotransmitter acetylcholine in synapse resulting continuous flow of neuromuscular signals causing paralysis and death of the insect (Kamanyire and Karalliedde, 2004; Singh and Rishi, 2005). Triazophos is the most indiscriminately used insecticide which affects the different non-target organisms like crabs, fish etc (Reddy et al., 1983). The bioaccumulation of this toxicant in earthworm affects seriously to the animals of its higher trophic levels (Darling and Thomas, 2005; Hobbelen et al., 2006; Van Gestel et al., 2011). It undergoes biomagnifications through food chain due to its persistence nature and causes serious problem to the human health (Naveed et al., 2011).

The earlier study indicates that the Triazophos is very toxic to earthworm *Eisenia fetida*. The median lethal concentration with contact filter paper test of Triazophos to the *Eisenia fetida* is the commencement of experiment. The commercial grade test chemical Triazophos (40% EC) was collected from the local market.

The Static replacement bioassays with the *Branchiura sowerbyi* was conducted in 500 ml Borosil glass beakers having 250 ml non-chlorinated tap water (APHA, 2012). The values of different physico-chemical parameters (temperature 25.0 ± 0.5 °C, pH 7.6

± 0.4 , free CO₂ 14.2 ± 0.8 mg/l, Dissolved Oxygen 4.9 ± 0.6 mg/l, total alkalinity 172 ± 5.6 mg/l as CaCO₃, hardness 117 ± 3.7 mg/l as CaCO₃) of test water during the experiment were estimated following the methods of APHA (2012). Each concentration of

Triazophos along with control used during acute toxicity test was accompanied by four replicates. Each replicate was provided with ten test organisms (*Branchiura sowerbyi*) with mean length of 11.6 ± 0.5 mm.

The rough range finding tests were initially performed to select the final test concentrations of Triazophos which were used to determine the median lethal concentration to *Branchiura sowerbyi*. The number of dead organisms was counted at every 24h of experiment. The dead organisms were removed immediately to avoid any organic decomposition. A certain amount of test water was replaced

every 24h by non-chlorinated stock water with a specific quantity of test chemical to make the desired concentration. Similar technique was also followed by the earlier

20

14.21 µg cm⁻² (Wang et al., 2011). The 1h EC

and EC90

values of

workers (Mukherjee & Saha, 2012; Bej et al., 2015). The median

Triazophos to zooplankton *Ceriodaphnia quadrangula* are 4.3

and 40.0 µg/l respectively. The 24, 48 and 72h LC50 values of Triazophos to *Cirrhinus mrigala* are 1.05, 0.87 and 0.75 mg/l respectively (Mahboob et al., 2015). The acute toxicity values of Triazophos to Crucian carp and common carp are 8.4 and 1.00 mg/l respectively (Ismail et al., 2009; Jin et al., 2010). The 96h LC50 value of Triazophos to *Anabas testudineus* is 0.270 mg/l (Jayakumar et al., 2014). There are no earlier reports on acute toxicity of Triazophos to benthic Oligochaetes. Therefore, the present study was aimed at evaluating the acute toxic effects of Triazophos to *B. sowerbyi* and their behavioural changes.

Materials and Methods

Test animal used in the bioassay was the benthic Oligochaete worm, *Branchiura sowerbyi* (Class: Oligochaeta, Family: Tubificidae). They were collected from local market. The test organisms were allowed to acclimatize in the test water for 72 hours before lethal concentration for 24, 48, 72 and 96h along with 95% con-

fidence limits were calculated with help of a statistical software programme (US EPA, 1999). The behavioural changes of the test organisms due to acute toxicity of Triazophos were also recorded during the bioassay.

The data of percent mortality of the test organisms was subjected to ANOVA using the R- software (R Development Core Team,

2011) followed by DMRT to determine significant variation among the means of control and treatments at different times of exposure.

Results and Discussion

No mortality of *Branchiura sowerbyi* was recorded in control during the experiment. The 24, 48, 72 and 96h LC50 values of Triazophos to *Branchiura sowerbyi* are given in Table 1. The mortal-

ity rate of *Branchiura sowerbyi* varied significantly ($p < 0.05$) with

RESEARCH 665

Volume : 5 | Issue : 5 | May 2016 • ISSN No 2277 - 8179 | IF : 3.508 | IC Value : 69.48

Research Paper

the increasing concentrations of the toxicant from 5.92, 5.60,

4.80 and 4.32 mg/l at 24, 48, 72 and 96h of exposure respectively (Table 2). A significant variation ($p < 0.05$) on the mortality rate was also recorded at each concentration (from 5.60 mg/l and above) with the increasing exposure time (24, 48, 72 and 96h). The clumping tendency and movement of *B. sowerbyi* were decreased with the increasing concentration of the test chemical and exposure time in comparison to control (Table 3). On the other hand, mucous secretion of the worm was increased with the increasing concentration and the time of exposure. The clumping tendency was totally absent at 7.20 mg/l and above test concentrations at all exposure times. But the rate of mucous secretion was very high at these concentrations.

In the present study, the 24, 48, 72 and 96 h LC50 values of Triazophos to *Branchiura sowerbyi* (8.17, 7.07, 5.81 and 5.04mg/l respectively) are comparatively much higher than the acute

toxic value of Triazophos to common carp (1.00 mg/l), *Anabas testudineus* (0.270 mg/l) and *Cirrhinus mrigala* (0.43-0.84 mg/l) (Ismail et al., 2009; Jayakumar et al., 2014; Mahboob et al., 2015). The 24 h LC50 value (8.17 mg/l) of Triazophos to *B. sowerbyi* corresponds with the LC50 value of crucian carp (8.4 mg/l).

The present experiment shows that the *Branchiura sowerbyi* is less susceptible to Triazophos than other aquatic animals. But it plays an important role in detritus food chain. The toxicant may be bioaccumulated in *Branchiura sowerbyi* which in turn may affect seriously to the animals of its higher trophic levels through food chain as was found in earthworms by the earlier workers (Darling and Thomas, 2005; Hobbelen et al., 2006; Van Gestel et al., 2011).

The values of median lethal concentrations of Triazophos to *Branchiura sowerbyi* will provide useful data for safe release of agricultural run-off containing Triazophos to the natural water bodies.

Table 1: LC50 values (with 95% confidence limits) of Triazophos to the *Branchiura sowerbyi* at different times of exposure (24, 48, 72 and 96h)

Test organ-ism	Concentration (mg/l)
24h	48h
<i>Branchiura sowerbyi</i>	8.17 (7.50-8.87)

Table 2: Mean values ($\pm$ SD) of *Branchiura sowerbyi* exposed to different concentrations of Triazophos at different times of exposure (24, 48, 72 and 96h). Mean values within columns indicated by different superscript letters (a-h) and mean values within rows indicated by different superscript letters (m-p) are significantly different (DMRT at 5% level)

Dose (mg/l)	% mortality of worm at different times of exposure (h)
-------------	--

24	48	72
00	00am $\pm$ 0.00	00am $\pm$ 0.00
4.0	00am $\pm$ 0.00	00 am $\pm$ 0.00
4.32	00am $\pm$ 0.00	00 am $\pm$ 0.00
4.80	00am $\pm$ 0.00	10 am $\pm$ 0.50
5.60	00am $\pm$ 0.00	20bn $\pm$ 0.71
5.92	10bm $\pm$ 0.00	30cn $\pm$ 0.50
6.40	20cm $\pm$ 0.00	40dn $\pm$ 0.50
7.20	30dm $\pm$ 0.00	50en $\pm$ 0.50
8.00	50em $\pm$ 0.83	70fn $\pm$ 0.71
8.80	50em $\pm$ 0.43	70fn $\pm$ 0.83
9.60	70fm $\pm$ 0.43	80gn $\pm$ 0.71
10.40	80gm $\pm$ 0.43	90hn $\pm$ 0.43
11.20	100hm $\pm$ 0.00	100hm $\pm$ 0.00

Table 3: Behavioural response of *Branchiura sowerbyi* (CT: clumping tendency; M: movement; MS: mucus secretion), –: absent, +: low, ++: moderate, +++: high) exposed to different concentrations of Triazophos at different times of exposure

Dose (mg/l)	Behavioural response of worm at different times of exposure
	24h

CT	M
0	+++
4.32	++
4.80	++
5.60	+
5.92	+
6.40	+
7.20	+
8.00	+
8.80	-
9.60	-

Acknowledgement

The authors are thankful to the Head, Postgraduate Department of Zoology and the Principal, Krishnagar Govt. College for providing infrastructural facilities to carry out the work.

CONCLUSION

The authors are thankful to the Head, Postgraduate Department of Zoology and the Principal, Krishnagar Govt. College for providing infrastructural facilities to carry out the work.

ref_str

1. American Public Health Association (APHA) (2012). Standard methods for the examination of water and wastewater. (Eds. Rice, E.W., Baird, R.B., Eaton, A.D.

& Clesceri, L.S.) American Public Health Association, American Water Works

Association, Water Environment Federation, Washington DC.

2. Bej, S., Mukherjee, D. and Saha, N.C. (2015). Acute toxicity of Alpha-Cyber- methrin to oligochaete worm, *Branchiura sowerbyi* (Beddard 1982) along with their behavioural responses. International Journal of Scientific Research.4(12), 27-28.
3. Darling, C.T.R., Thomas, V.G. (2005). Lead bioaccumulation in earth worms, *Lumbricus terrestris*, from exposure to lead compounds of differing solubility. Sci.Total.Environ. 346, 70–80.
4. Hart, K. & Pimentel, D. (2002). Public health and cost of pesticides. In: *Encyclopedia of pest management* (ed. D. Pimentel), Marcel Dekker, New York, pp.677–679.
5. Hobbelen, P.H.F., Koolhaas, J.E., van Gestel, C.A.M. (2006). Bioaccumulation of heavy metals in the earthworms *Lumbricus rubellus* and *Aporrectodea caliginosa* relation to total & available metal concentrations in field soils. Environ. Pollut. 144, 639–646.
6. Ismail, M., Ali, R., Waheed, U. & Khan, Q. M. (2009). Evaluation of the acute toxicity of profenofos & its effects on the behavioral pattern of fingerling common carp (*Cyprinus carpio* L., 1758). *Bull. environ. Contam. Toxicol.* **82**, 569–573.
8. Jayakumar, D. Sudarsanam, Garg Ramesh, P. Praveena (2014). Evaluation of Tri- azophos Induced Histopathology and Recovery in a Fish *Anabas testudineus*.
9. International Journal of Toxicological and Pharmacological Research 2014.6(4), 67-74.
10. Jain S, Mythily S, Ahmed RS, Arora VK, Banerjee BD. (2010). Induction of oxidative stress and histopathological changes by sub-chronic doses of triazophos. Indian J Biochem Biophys 47(6), 388-92.
11. Jin, L., Ravella, R., Ketchum, B., Paul, R.,
12. Bierman, R., Heaney, P., White, T. And
13. Brantley, S. L. (2010). Mineral weathering & elemental transport during hillslope evolution at the susquehanna/shale hills critical zone observatory. *Geochim.Cosmochim.Acta*, **74**, 36693691.
14. Kamanyire R, Karalliedde L. (2004) Organophosphate toxicity & occupational exposure. Occup Med (Lond). 54(2), 69-75.
15. Mahboob, S., Ghazala, S., Sultana, S., Al-Akel, A.S., Humood, F. A.B., Al-Misned, F. and Zubair, M. (2011). Pesticide residues in flesh of *Cirrhinus mrigala* collected from a commercial farm & river Chenab at Trimu Head, Jhang. *Pakistan J.Zool.* **43**, 97-101.
16. Mingjing Q, Zhaojun H, Xinjun X, Lina Y. (2003). Triazophos resistance mechanisms in the rice stem borer (*Chilo suppressalis* Walker). *Pestic Biochem Physiol.* 77(3), 99-105.
17. Mukherjee, D., and N.C. Saha. (2012). Acute toxicity of 2,4,6-Trichlorophenol to copepod, *Cyclops viridis* and oligochaete worm, *Branchiura sowerbyi*. *Environment and Ecology.* 30(3C), 1165–1170.
18. Naveed, P. Venkaeswarlu and C. Janaiah (2011). Biochemical alteration in-666 IJSR - INTERNATIONAL JOURNAL OF SCIENTIFIC RESEARCH

Research Paper

Volume : 5 | Issue : 5 | May 2016 • ISSN No 2277 - 8179 | IF : 3.508 | IC Value : 69.48

duced by triazophos in the blood plasma of fish, *Channa punctatus* (Bloch). *Annals of Biological Research.* 2 (4), 31-37.

19. Reddy S P, Bhagyalaxmi A, Rammurthi R. (1983). *Toxicol Lett* 22, 335-338.
20. R Development Core Team (2011). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL <http://www.Rproject.org/>. Accessed 8 Oct 2012.

21. Singh M, Rishi S. (2005). Plasma acetylcholinesterase as a biomarker of triazophos neurotoxicity in young and adult rats. *Environ Toxicol. Pharmacol.* 19(3), 471-6.

22. US EPA.(1999). Probit program version 1.5. Ecological Monitoring Research Division, Environmental Monitoring Systems Laboratory, US Environmental Protection Agency, Cincinnati, Ohio 45268. <http://www.epa.gov/nerleerd/stat2.htm>. Accessed 14 Jan 2012.
23. Van Gestel, C.A.M., Ortiz, M.D., Borgman, E., Verweij, R.A. (2011). The bioaccumulation of Molybdenum in the earthworm *Eisenia andrei* influence of soil properties and ageing. *Chemosphere*. 82, 1614–1619.
24. Wang, Yanhua, Tao Cang, Xueping Zhaon, Ruixian Yu, Liping Chen, Changxing Wu, Qiang Wang. (2012). Comparative acute toxicity of twenty-four insecticides to earthworm, *Eisenia fetida*, *Ecotoxicol. Environ. Saf.* doi:10.1016/j.ecoenv.2011.12.016.
25. Worthing, C.R. and R.J. Hanrce. (1991). The pesticide manual: a world compendium, ninth ed., *British Crop Protection Council*, Surrey, pp. 83.



IJSURP Publishing Academy

International Journal Of Scientific And University Research Publication
Multi-Subject Journal

Editor.

International Journal Of Scientific And University Research Publication



+965 99549511



+90 5374545296



+961 03236496



+44 (0)203 197 6676

www.ijsurp.com