



International Journal Of Scientific And University Research Publication

ISSN No **2364/2017**

Listed & Index with
ISSN Directory, Paris



Multi-Subject Journal



ALLELOPATHIC ACTIVITIES OF SPECIFIC MICROBIAL METABOLITES IN THE INLAND PRAWN FISHERIES OFF EASTERN UTTAR PRADESH, INDIA

Sushil K. Upadhyay || Zoology Department
Science Faculty
Swami Vivekanand Subharti University
Meerut-250005
Uttar Pradesh
India

The gram positive bacterial strain of *Streptococcus* spp. recovered from *Macrobrachium rosenbergii* pond water was found to be a pathogenic strain. The in vitro allelopathic activities of all 22 bacterial

strains of same habitat were tested against *Streptococcus* spp. Out of aforementioned strains only one showed antagonistic activity and identified as gram positive rod, *Clostridium* spp. The in vitro allelopathic interactions of these two bacterial strains were demonstrated to be a potential antagonist in the inland water fisheries. This study was carried out for the determination of natural allelopathic and antibacterial activity of antagonistic bacterial strain against the pathogenic bacterial strains of fresh water aquaculture system of eastern Uttar Pradesh, India.

Macrobrachium rosenbergii
Allelopathic activity, Antibacterial

مقدمة

Aforementioned applications of approaches are using live bio-therapeutic agents such as bacterial isolates and metabolites to be secreted by them present in the cell free supernatant. The antiseptic strains in their *Lacto bacillus* bacteriocins are produced by several secretory metabolites have been used against the gram positive food *Staphylococcus*, *Listeria monocytogenes* borne pathogenic strain of *Clostridium perfringens* and spores of *Bacillus subtilis*, *aureus*. Therefore, now a days microbial antagonists with allelopathic bustle may concerned much attention for the use as native and natural biopreservatives in food and fisheries industries (Saranya and Hemashenpagam, 2011; Kaynar and Beyatli, 2012). However, studies relating to the antibacterial properties of these organisms have been limited and not fully exploited for use. In this study we report the benefits of carefully selected strain of gram positive spp. on the growth performance of giant fresh *Clostridium* bacilli, as biological control agent *Macrobrachium rosenbergii* water prawn, in aquaculture by their allelopathic activity against the pathogenic spp. Such a strategy could well form the *Streptococcus* bacterial strain, basis of a rationale for the search of potential antagonistic and allelopathic antibacterial agents that could help to bring down disease incidence in the scampi culture pond and inland water fisheries.

Materials and Methods

The bacterial samples were collected from the water and soil of the pond which was being cultured in *Macrobrachium rosenbergii* pond in which Allahabad, U.P. Simultaneously bacterial sample were collected to by swabbing *M. rosenbergii* body surface of giant freshwater prawn. Bacterial samples were cultured in nutrient broth for 24hrs and 0.1ml of culture aseptically inoculated in nutrient-agar. The pure culture of each isolates was created, serially diluted and colonies obtained were counted after Upadhyay, Malhotra and Malhotra (2009). Antagonism of bacterial strain 158 against target microbial culture 144 was detected by disc diffusion assay (Chang, Liu and Shyu, 2000). The sterilized discs of Whatmann paper were placed aseptically on cultured pathogenic susceptible bacteria, previously soaked with cell free broth culture of test organism (antagonist) followed by incubation of culture at 36°C and periodically examined for the zone of clearance around the disc in cultured medium as a positive indication of inhibitory activity. The growth pattern of target strains was calculated after every 3hrs for 58hrs by The growth curve was plotted, spectrophotometer (ELICO) at 600nm along the diameters of the zone of clearance.

Results

The samples of bacterial pure cultures isolated from scampi, pond water and soil were subjected to different biochemical tests for their numeric evaluation and preliminary identification. An antagonism

The enhanced demand for aquatic food products has imposed development and strengthening of aquaculture activity in most regions of the world. This intensification has led to an increased use and misuse of drugs and chemicals in aquaculture systems that lead to emergence of food safety concern. Microbial products are seen as, among other things, alternatives to the prophylactic use of chemicals (Moriarty, Decamp, and Lavens, 2005). However the poor qualities of probiotics in the market and the misapprehensions regarding their function have raised doubts on their efficacy. The enhanced use of antibiotics and other chemical compounds may illicit and stimulate as leads to the resistance among pathogenic bacterial strains internally posing a great challenge to human beings (Jebasingh and Murugan, 2011). The inhibitory effect of biochemical compounds against the other flora and fauna is called allelopathy. The word allelopathy derived from two Greek words *allel* and *pathos* and uttering the meaning as "let the one suffer that looks like you". In other terms we can say that allelopathy is the direct or indirect interactive impact of the biochemical metabolites of microbial strains against the others that escape into the environment. In aquatic systems the definition of allelopathy so broad and comprises all chemical interactions among unicellular organisms (Cembella, 2003; Gross, 2003; Legrand, Rengefors, Fistarol, and Granéli, 2003; Granéli, 2006; Ianora, Boersma, Casotti, Fontana, Harder, Hoffmann, Pavia, Potin, Poulet, and Toth, 2006; Tillmann, John, and Cembella, 2007). At present forage is no longer taken in consideration by the heterotrophs only for taste and instantaneous nutritional requirements for survival, but also in terms of their utility of give specific health benefits beyond their fundamental nutritive quality. Now a days gross part of the active running food markets are influenced by the healthy food products aiming towards up healing the balance between micro flora of gut and surrounding mediums. The antagonistic possessions of bacterial strains can be employed in various systems like aquaculture, agriculture, and epidemiology, biological control of pests and in eradication of disease causing or pathogenic organisms (flora and fauna). The pattern and specificity of the allelopathy with the corresponding pathogenic susceptible strains is based on the lowering of pH values, inhibiting properties of their metabolites, including lactic acid, hydrogen peroxides and bacteriocidin and subsidizing to the inhibition of pathogenic bacterial strains has been worked out by the earlier workers (Servin, 2004; Ligockaa, Paluszak and Hermann, 2005; Vuyst and Leroy, 2007; Szala, Paluszak, and Motyl, 2012). Decent and innovative approaches have been used since yesterdecades as alternative to antibiotics in control and treatment of several gastrointestinal, infectious, microbial and parasitic diseases.

of inhibition increased with enhanced level of antagonist. Therefore, as a potential probiotic, strain 158 has to be supplied in sufficient quantity and on a regular basis into the system. One of the most viable criteria for considering the bacterium as a candidate to be used in biocontrol program in prawn culture system is its non-pathogenicity (Verschuere et al., 2000). Culture 158 has not been recognized and reported as prawn pathogen previously. The present work strengthens the candidature of strain 158 as an “antagonistic/allelopathic probiotic” in prawn rearing systems.

ref_str

- (2003). Chemical ecology of eukaryotic microalgae **Cembella, A.D.** .1 in marine ecosystems. *Phycologia*, 42(4), 420-447.
- , and Shyu, C. Z., (2000). Use of prawn blood **Chang, C.I., Liu, W.** .2 agar hemolysis to screen for bacteria pathogenic to cultured tiger prawns *Pennaeus monodon*. *Diseases in Aquatic Organism*, 43, 153-157.
- (2006). Kill your enemies and eat them with the help of **Granéli, E.** .3 your toxins: an algal strategy. *African Journal of Marine Science*, 28(2), 331-336.
- (2003). Allelopathy of aquatic autotrophs. *Critical Reviews in Plant Sciences*, 22(3-4), 313-339.
- , Casotti, R., Fontana, A., Harder, J., **Ianora, A., Boersma, M.** .5 Hoffmann, F., Pavia, H., Potin, P., Poulet, S.A., and Toth, G., (2006). New trends in marine chemical ecology. *Estuaries and Coasts*, 29(4), 531-551.
- , and Murugan A. (2011). Antagonistic activity of **Jebasingh S.E.J.** .6 the Barnacle (*Balanus amphitrite*) associated bacteria against human bacterial pathogen. *World Journal of Medical Sciences*, 6(1), 36-41.
- and Beyatli, Y. (2012). Antagonistic Activities of **Kaynar, P.** .7 *Bacillus* spp. Strains Isolated from the Fishes. *Journal of Applied Biological Sciences*, 6(3), 77-81.
- K., Fistarol, G.O., and Granéli, E., 2003. **Legrand, C., Rengefors, S.** .8 Allelopathy in phytoplankton-biochemical, ecological and evolutionary aspects. *Phycologia*, 42(4), 406-419.
- ., and Hermann, J. (2005). Influence of **Ligockaa, A., Paluszak, Z.** .9 physico-chemical factors on the effectiveness of bacteriocins of lactic acid bacteria activity against the pathogens in sewage sludge. *Mediterranean Wetlands*, 61, 413.
- , and Lavens, P. (2005). Probiotics in **Moriarty, D.J.W., Decamp, O.** .10 *Aquaculture*, *Aquaculture Asia Magazine*, p14-16.
- and Hemashenpagam, N. (2011). Antagonistic activity **Saranya, S.** .11 and antibiotic sensitivity of Lactic acid bacteria from fermented dairy products. *Advances in Applied Science Research*, 2(4), 528-534.
- . (2004). Antagonistic activities of lactobacilli and **Servin, A.L.** .12 bifidobacteria against microbial pathogens. *FEMS Microbiology Review*, 28, 405.
- Paluszak, Z., and Motyl, I. (2012). Antagonistic effect of **Szala, B.** .13 lactic acid bacteria on *Salmonella* senftenberg in mixed culture. *Poland Journal of Environmental Study*, 21(5), 1399-1403.
- , John, U., and Cembella, A., (2007). On the **Tillmann, U.** .14 allelochemical potency of the marine dinoflagellate *Alexandrium ostensfeldii* against heterotrophic and autotrophic protists. *Journal of Plankton Research*, 29(6), 527-543.
- Malhotra, A., and Malhotra, S.K. (2009). **Upadhyay, S.K.** .15 Antagonistic microbial interactions with specific metabolites activity patterns in aquatic and terrestrial host parasite systems. 21st National Congress Parasitology, p60.
- , Rombout, G., Sorgeloos, P., and Verstraete, W. **Verschuere, L.** .16 (2000). Probiotic bacteria as biological control agents in aquaculture. *Microbial Molecular Biology Review*, 64(4), 655-671.
- and Leroy, F. (2007). Bacteriocins from lactic acid **Vuyt, L.** .17 bacteria. Production, purification and food application. *Journal of Molecular Microbiology and Biotechnology*, 13(4), 194-199.

assay by disc diffusion showed that cell free supernatant of sample 158 inhibited growth of sample 144 with zone of clearance ranged between 0.6 to 2.1cm in diameter (Fig.1). No growth of antagonistic bacteria was commenced between 0.0 to 4.0hrs (Fig. 2). The exponential growth was take place from 6 to 42hrs as assessed by the rapid change in optical density (OD). However, very little change was noticed in OD516 IJSR - INTERNATIONAL JOURNAL OF SCIENTIFIC RESEARCH between 42 to 58hrs. This was the stationary phase. In this phase maximum antagonistic activity was observed at 38hrs (Fig.2). Maximum zone of clearance (2.1cm) was obtained at 38hrs and minimum obtained at 0.0hrs.

The colonies were smooth, circular, with regular **Strain 158:** edges, convex, shiny and mucoid. The rate of growth was very good. After 24hrs, culture did not settle down in broth but caused turbidity. These were gram positive rods with endospores, motile and isolated from fresh water scampi pond's water of selected sites. These were spp. *Clostridium* identified as

The colonies were evoked blue in color, rough, dry and **Strain 144:** formed spores. The growth on nutrient agar was extremely slow and optimum growth commencing after 72hrs of incubation. Very little or no growth was recorded in nutrient broth. Gram staining revealed positive streptococci. The strains i.e. these to be the chains of cocci and identified *M. rosenbergii* were recovered from body surface of *Streptococcus viridians* as

niosucDis

Since the first antibiotic from bacterium was described in 1966, the number of new compounds has increased constantly during the years. Even though only a few compounds from organisms might be interesting for the pharmaceutical market today. Some bacterial species are already used as biocontrol, and are added to aquaculture stocks. Most metabolites from bacteria found so far were isolated from marine bacterial species. In recent studies wide array of bacteria were tested for bacterial antagonistic effects, and it was demonstrated that this trait appeared to be a widespread feature in marine habitats, and was present in many bacterial groups. The production of secondary metabolites by bacteria, has been reported previously, and some organism *Roseobacter* namely, of this group is thought to be either probiotic or antibiotic in different aquaculture conditions. The bacterial strain 158 inhibited the growth of other strain 144 when the initial count of the antagonist i.e. 158 was far greater than that of the prawn pathogen, 144. This indicated that the antagonist must be available at significantly higher levels than the pathogen, and the degree of inhibition increased with enhanced level of antagonist. Therefore, as a potential probiotic, strain 158 has to be supplied in sufficient quantity and on a regular basis into the system. One of the most viable criteria for considering the bacterium as a candidate to be used in biocontrol program in prawn culture system is its non-pathogenicity (Verschuere et al., 2000). Culture 158 has not been recognized and reported as prawn pathogen previously. The present work strengthens the candidature of strain 158 as an “antagonistic/allelopathic probiotic” in prawn rearing systems.

sentemgdelownkCA

SKU is grateful to Prof. Jayshree Naidu, Department of Microbiology, Modern Science P.G. College (autonomous), Jabalpur, M.P. for her help in identification of bacterial strains.

.This indicated that the antagonist must be available at significantly higher levels than the pathogen, and the degree



IJSURP Publishing Academy

International Journal Of Scientific And University Research Publication
Multi-Subject Journal

Editor.

International Journal Of Scientific And University Research Publication



+965 99549511



+90 5374545296



+961 03236496



+44 (0)203 197 6676

www.ijsurp.com