



International Journal Of Scientific And University Research Publication

ISSN No **2364/2018**

Listed & Index with
ISSN Directory, Paris



Multi-Subject Journal



COMMUNITY DYNAMICS AND FLORISTIC COMPOSITION OF NATURAL VEGETATION IN ABAYA HAMASSA, RIFT VALLEY OF ETHIOPIA

Eyasu Chama Didana || Department of Biology
Faculty of Natural and Computational Science
Wolaita Sodo University
Po Box 138
Wolaita Sodo
Ethiopia.

Ethiopia, with its multidimensional biogeo terrain is unique for its species diversity and bizarre dispersion of wood land. A total of 315 plant species representing 198 genera

ts species diversity and bizarre dispersion of wood land. A total of 315 plant species representing 198 genera and 59 families were identified in Abaya-Hamassa natural vegetation in the rift valley of Ethiopia. Six plant communities were identified and described. Community-environment correlation was assessed based on Ecological, physiological and pedological data collected from 55 quadrates (400 square meters) belonging to eight selected sites between October 2013 to June 2014. The environmental parameters such as altitude, slope, aspect, PH, electrical conductivity, total nitrogen and soil texture (% sand, % clay and % slit) were measured for all quadrats. Among the environment parameters, altitude was negatively correlated with sand and clay. The floristic composition, floral diversity, identification of community type and community-environment interaction were investigated. The Percentage of coverage, Canopy, abundance and frequency of species were estimated and converted as per the modified Brawn-Blanquet 1-9 scale.

Abaya Hamassa, Biodiversity,الكلمات الرئيسية
 Communit,Environment ,interaction,

crucial for proper management. This is due to the fact that knowledge about vegetation composition and structure provides insight about the potential of plant community to succeed after disturbance.

1-Description of the Study Area

The study area is located in between Wolaita and Gamo-Gofa zones at 6 27.404°N - 637.368°N and 6 44.042°E - 3652.169° E, in the Southern Nations, Nationalities and Peoples' Regional State (SNNPRS) within the central Rift Valley of Ethiopia. The area is located to the east of western escarpment of Rift Valley to the west and northeast of Lake Abaya. The elevation ranges from 1184 m.a.s.l. at the coast of Lake Abaya to 1410 m.a.s.l. at Wanke derya. The elevation decreases towards the Lake by forming tiers in a long distance. The drain- age pattern follows the general topographic orientation from west to east. Small rivers and streams rising from Wolayta and Chenchha highlands drain to Lake Abaya and Lake Chamo. The soil along the floor of the Rift Valley and Lake Abaya are alluvial of Hare stream (Vaukasinvic, 1969) and colluvial materials and lacustrine deposits of the Pleistocene (Mohr, 1971) categorized also as Fluvisols of the FAO/ UNESCO revised legend (1990). The average annual rainfall of the study area ranges from 650 to 800 mm. Rainfall is bimodal occurring from March to may (short rain) and June to September (long rain). The mean monthly temperature of the area is between 22 to 28°C (Wolayta Zone Agricultural Office, 2013).

2-Vegetation

The vegetation types of semi-arid regions in Ethiopia woodland, bush-land and tickets, types belonging to *Acacia* include open grasslands (Coppock, 1994). White (1983) classified the vegetation of Ethiopia under the Sudanian Regional Center of Endemism, Somalia-Massai Regional Center of Endemism, Afromontane Regional Center of Endemism and Afro Alpine Archipelago-like Regional Center of Endemism. The present study area can be plunged in the Somalia-Massai Regional Center of Endemism but if based on the broad categorization of the vegetation, the study area was mainly small-leaved deciduous woodland type. woodland was found mainly *Acacia-commiphora* Some sort of between altitudes of 500-1900 m. a. s. l. with an average annual temperature of 18°C to 27°C and rainfall between about 410-820 mm. This woodland area had traditionally been a grazing land (1992).et al., (Ensermu Kelbessa

مقدمة

Biodiversity refers to the variety of life forms, genetic diversity and the assemblage they form. According to Kumer (1981), and Paul (1993) species diversity can be viewed in terms of species richness, species endemism, evenness and taxonomic diversity. Biological systems, whether tundra, forests, savannahs, grasslands, deserts, lakes, rivers, wetlands, coastal communities or marine ecosystems are functionally complex and this complexity is associated in often obscure ways, with the diversity of their component species (U.S. National Research Council, 1992). Most of the estimates suggest that there are about 250,000 species of vascular plants in the world (U.S. National Research Council, 1992). Approximately two-thirds of these are found in the tropics. One-sixth of the earth's diversity of plant life about 45,000 species can be found in Latin America followed by Africa. The diversity of higher plant species increases as one moves from poles to the equator. Ensermu Kelbessa estimated about 6000 species of higher plants in Ethiopia, of which nearly 10.5% are endemic. The alarming population increase obviously leads to an increase in the demand for natural resources, including arable land, water, wood for construction and energy. The increase in population size, urban development, and expansion of commercial and subsistence farming has exerted considerable influence on woodland and vegetation in the Rift Valley region of Ethiopia. For the last few decades much of the natural vegetation had been converted into crop fields for undertaking intensive farming activities. Deforestation for various domestic exercises, demand for charcoal had decimated the vegetation. Most of the land is now either cultivated or barren leaving the woodland fragmented into small patches (Zerihun .. 1999). It seems thus doubtful that the forests once et al/Woldu, covered 40% of the country (FAO,1980) had reduced to negligible percentage today, with only 2.7% of the country extremely forested *Podocarpus* and 0.9% of the original Coniferous forest (forests and 7.6% of the *Junperus*), 11% of the original broad leaved original savanna woodlands remained depicting an alarming picture for future generations (WCMC,1991). The Rift Valley of Ethiopia is generally characterized by semiarid climatic condition. Although there subsist a number of wood- land and bush land resources but literature on these vegetation is meager. The natural vegetation of Abaya-Hamassa is under threat by the combined forces of resources exploitation and for-est degradation due to activities akin to frequent burning, clearing and grazing. In order to suggest the mechanisms to maintain and restore these resources on a sustainable basis, a clear understanding on vegetation structure and floristic composition is

the process of digestion organic nitrogen would be converted into ammonium nitrogen. Nitrogen was estimated then from the amount of ammonia liberated by distillation of the digestion with alkali. The procedure started by taking one gram of air dried soil that had passed through 0.5mm sieve. It was placed in a dry 500ml Kjeldahl flask, mixed with 7g of potassium sulfate and 0.8g of copper sulfate to catalyze the reaction and 12ml of concentrated sulfuric acid was added to it. The digestion flasks were then put to the digestion system which was pre-heated (420 deg. Celsius). Samples were allowed to be digested for about one hour until all the samples were clear of green colour. After the racks of the tubes were cooled for 15 minutes, 75ml of distilled water was added into the tube. In a conical flask 25ml of receiver solution was added and the flask was placed into the distillation unit. In the distillation unit 50ml of 40% sodium hydroxide was dispensed and distillation allowed to go for about 4 minutes until the receiver solution in the distillate flask became green in colour indicating the presence of alkali ammonia. The distillate was then titrated with standardized 0.1N HCl until the blue gray end point was reached. The whole reaction can be summarized as follows:- K_2SO_4

THE EQUATION IS NOT IN A STRAIGHT LINE, **5-Digestion:** SHOULD BE WRITTEN IN A SINGLE LINE.

6-Distillation:

The percentage of total nitrogen was then calculated as follows:-

$$\text{Titration of volume for sample } T = \frac{\text{Titration volume for blank } B}{\text{Titration volume for blank } B}$$

To measure soil electrical conductivity a soil *normality of acid* $N =$ water suspension in a ratio 1: 2.5 was prepared. In this case soil water suspension was made by dissolving 10g of soil by 25ml of distilled water and stirred for 30 minutes with automatic stirrer. Then electrical conductivity was measured using conductivity meter. All the environmental data is presented in Table-6 and 7.

7-Data analysis

Multivariate techniques were used to study the complex nature of plant communities. In aiming for a detailed floristic vegetation description, two major approaches were distinguished. These were the relevé analysis for classification and the continuum analysis for ordination (Mueller-Dombois and Ellenberg, 1974). These two approaches toward vegetation study were based on different concepts of the essential nature of vegetation. Classification aims at grouping individual stands into categories. The stands those were similar to one another form one class, which was separated from other classes that also consist of similar stand (Mueller-Dombois and Ellenberg, 1974). This method emanated from the belief that vegetation was composed of certain distinct and fairly discrete plant communities (the concept of community unit theory) (Whittaker, 1962 and 1967; Shimwell, 1971). The properties common to a group of similar stands in a class were then abstracted to serve as description of that class. Plots were grouped into clusters with aid of the program TWINS-SPAN. Jaccard's coefficient of similarity was computed to see the species composition turnover of the communities and species composition grades gradually or in discontinuity (Table-4). Sørensen's Coefficient was calculated to analyze the species similarities between the study areas with other natural vegetation types in different parts of the country (table-8).

8-Description of floristic composition

A total of 315 species of plants representing 198 genera 59 families were recorded from the study area were summarized in the Table-I. Of these, Poaceae had 39 (12.38%) species standing largest and some 19 families had each 1 (0.32%) species as least. Out of these species identified in the study area, 145 (45.63%) species were herbs, 76 (24.06%) species were shrubs, 38 (12.19%) species were trees, 36

3-Vegetation Data Collection

In order to identify sampling sites reconnaissance survey was made and eight sampling sites were selected based on physiognomy of the vegetation. Then sample plots were selected and vegetation data was collected from all the 55 quadrats. During sampling, visually checked homogeneous representative stands were selected and delimited for sampling. At each sampling area, plots (quadrats) of 400 m² (20 m x 20 m) were measured. Inside quadrat, nested plots of 10 m x 10 m for trees, 5 m x 5 m for shrubs and 2 m x 2 m for grasses and other herbs were established. All plant species in the quadrats were recorded and voucher specimens were pressed in the field for subsequent identification at the Herbarium of Wolayta Sodo University. Percentage cover values of trees, shrubs, herbs and grasses were estimated in the field and later converted to cover/abundance values of Braun-Blanquet scales as modified by van der Maarel (1979). The collected specimens were identified by using standard keys as published one to nine Volumes of Flora of Ethiopia and Eritrea.

4-Environmental Data

The following environmental data and topographic parameters were gathered for each plot. Altitude, slope, aspect and position for each plot were determined respectively with an Altimeter, Suunto Clinometer, Suunto compass and GPS-48. Soil samples were collected from 5 sites in each plot (quadrat), four at the corners and one at the middle of the plot at 0-20 cm. Composite soil samples weighing about 1.5 kg were brought to the soil laboratory for further analysis. The samples were air dried, sieved with a mesh size of 0.5 mm and 2 mm, were analyzed at the soil laboratory, following the methods of Jackson (1973) and Juol (1978). The analyzed parameters were soil texture, pH, total nitrogen and electrical conductivity. Texture was determined using Bouyoucos Hydrometer method; pH was measured using a pH meter at 1:2.5 distilled water to soil suspension while total nitrogen was determined by using Kjeldahl method. Electrical conductivity was measured by using conductivity meter (Table-6). Fifty grams of 2 mm sieved dried soil sample was added to 50 ml of 5% sodium hexametaphosphate along with 100 ml distilled water. The suspension was stirred with a glass rod and then allowed to stand for 30 minutes. The soil suspension was stirred using a multi-mix machine for 15 minutes, and then transferred into a one liter glass cylinder. The suspension was diluted with distilled water up to a one liter mark. The soil suspension was mixed by covering the mouth of the cylinder by hand and inverting it several times. Hydrometer and temperature readings were taken at 40 seconds and the soil suspension was allowed to settle for 2 hours. After 2 hours the readings of Hydrometer and temperature were taken for the second time. Then the different soil fractions were calculated as follows:

$$\% \text{ SAND} = 100 - [H_1 + 0.2 (T_1 - 68) - 2] \times 2.1$$

$$\% \text{ CLAY} = [H_2 + 0.2 (T_2 - 68) - 2] \times 2.2$$

$$\% \text{ SILT} = 100 - (\% \text{ SAND} - \% \text{ CLAY}) \times 3$$

Where H_1 = the first hydrometer reading after 40 seconds.

T_1 = the temperature reading after 40 seconds. H_2 = the hydrometer reading after 3 hours.

T_2 = the temperature reading after 3 hours. Soil pH was determined following Juol (1978). A 1: 2.5 soil suspension in water was made by dissolving 10g of soil that has passed through 2 mm sieve with 25 ml of distilled water. The suspension was stirred with a glass rod and allowed to settle for 30 minutes. The pH was measured using pH meter. Nitrogen was determined using the macro Kjeldahl method. The analysis of soil nitrogen requires a complete breakdown of the organic nitrogen (digestion), followed by distillation and titration. In

	e		
0.32	1	Ochnaceae	50
0.32	1	Onagraceae	51
0.32	1	Plumbaginaceae	52
0.32	1	Portulacaceae	53
0.32	1	Pteridaceae	54
0.32	1	Salvadoraceae	55
0.32	1	Santalaceae	56
0.32	1	Scrophulariaceae	57
0.32	1	Sinaroubaceae	58
0.32	1	Zygophyllaceae	59

Based on the TWINSpan output and ecological evaluations in field, six ecologically meaningful community types (clusters) designated as I, II, III, IV, V and VI were identified. The plant communities were named by the dominant and/or characteristic species, which occur in each group, using the relative magnitude of mean cover/abundance. The mean cover abundance of major species in the community type was estimated. As a result, the six community types were named after two or three of the dominant and/or characteristic species and the communities are described as given below (table-2).

9-Communities

Canthium setiflorum – *Acacia hockii* – *Pappea capensis* Community I.

This community is found at an altitudinal range of 1184–1309 are *Combretum molle* and *Pappea capensis*, *Acacia hockii*, m.a.s.l. *Rhus natalensis*, dominant tree species. The shrub layers are *Less Ximenia americana*, *Acacia brevispica* and *Grewia bicolor*. *Commiphora* abundant but still dominant tree species are *Africana*, *Terminalia brownii* and *Boswellia rivaie*. *Hyparrhenia rufa*, *Cynodon dactylon*, *Laggera crispata*, *Heteropogon contortus*, are elements of the *Bothriochloa insculpta*, and *Abutilon firarianum* *Cardiospermum halicacabum*, *cissuss* herb layer of the community. *quadrangularis*, *C. rotundifolia*, *Cyphosemma adenocaula* and are climbers encountered in this community. *kleinia squarrosa*

Commiphora africana – *Acacia Senegal* – *Balanites aegyptiaca* types. Community II

The altitudinal range of this community is from 1241-1410m.a.s.l. The dominant tree-shrub species *fricana*, *Rhus natalensis*, *Acacia senegal*, *Harrisonia* a *Commiphora* are *abyssinica*, *Balanites aegyptiaca*, *Ximenia americana*, *Dichrostachys Grewia* and *cineria*, *Mystrxylo aethiopicum*, *Ziziphus mucronata*, *villosa*. *Helichrysum schiperi*, *Jastica flava*, *Leptochloa rupestris*, *Sehima nervosum*, *Cyperus dubiud*, *Acalypha ciliate*, *Clitoria ternatea*, herb surface layers. *Sansevieria ehrenbergii*

inca. *Zanthoxylum chltbeum* - *Commiphora habess* Community III types.– *Grewia villosa*

This community type is found at an altitudinal range of 1233– is characteristic shrub species *Zanthoxylum chalybeum*, 1283masl. *Terminalia brownie*, *Commiphora habessinica*, of the community. *Boswellia riae*, *Mystrxylon aeyhiopicum*, *Grewia villosa*, *Croton zambesicus*, *Diospyros abyssinica*, *Grewia velutina*, *Tamarindus Leptochloa rupestris*, are dominant tree – shrub species. *indica* are dominant grass species in *S. ioclados* and *Sporobolus pyramidals*, the community.

Acalypha fruticosa – *teclea nobilis* – *Baphia abyssinica* Community IV

(11.25%) species were trees/shrubs, 18 (5.63%) species were climber and 2 (0.63%) species were ferns and epiphytes each. Species such *Acacia brevispica*, *Acacia tortilis*, *Acalypha fruticosa*, *Baphia abyssinica*, *Cissus quadragularis*, *Combretum molle*, *Croton zambesicus*, *cinerea*, *Diospyros abyssinica*, *Euphorbia tirucalli*, *Grewia bicolor*, *Grewia velutina*, *Harrisonia Dichrostachys abyssinica*, *Lecaniodiscus fraxinifolius*, *Maerua triphylla*, *Pap-pa capensis*, *Rhus natalensis*, *Sansevieria abyssinica*, *Sansevieria ehrenbergii*, *Tamarindus indica*, *Teclea nobilis*, *Terminalia brownii*, *Ximenia* were widely distributed species *Ziziphus mucronata* and *americana*, with high frequency.

Percent (%) of composition	No. of species recorded	Family	Serial No
12.38	39	Poaceae	1
12.06	38	Fabaceae	2
6.98	22	Asteraceae	3
4.76	15	Euphorbiaceae	4
4.44	14	Malvaceae	5
4.13	13	Acanthaceae	6
3.81	12	Tiliaceae	7
4.49	11	Cyperraceae	8
3.17	10	Amaranthaceae	9
2.86	9	Lamiaceae	10
2.22	7	Rubiaceae	11
2.22	7	Vitavaceae	12
1.90	6	Celastraceae	13
1.90	6	Cappariaceae	14
1.90	6	Asclepiadaceae	15
1.59	5	Spindaceae	16
1.59	5	Anacardiaceae	17
1.59	5	Moraceae	18
1.59	5	Solanaceae	19
1.59	5	Burseraceae	20
1.27	4	Polygonaceae	21
1.27	4	Asparagaceae	22
1.27	4	Commelinaceae	23
1.27	4	Cucurbitaceae	24
1.27	4	Oleaceae	25
0.95	3	Boragintaceae	26
0.95	3	Chenopodiaceae	27
0.95	3	Combretaceae	28
0.95	3	Crassulaceae	29
0.95	3	Dracaenaceae	30
0.95	3	Rhaminaceae	31
0.63	2	Apiaceae	32
0.63	2	Flacourtiaceae	33
0.63	2	Annonaceae	34
0.63	2	Apocynaceae	35
0.63	2	Balanitaceae	36
0.63	2	Ebenaceae	37
0.63	2	Loranthaceae	38
0.63	2	Olacaceae	39
0.63	2	Rutaceae	40

Table 1. The summary of floristic composition identified in the study area.

0.32	1	Actinopterida cea	41
0.32	1	Aizoaceae	42
0.32	1	Aloaceae	43
0.32	1	Anthericaceae	44
0.32	1	Cactaceae	45
0.32	1	Loganiaceae	46
0.32	1	Molluginaceae	47
0.32	1	Myrtaceae	48
0.32	1	Nyctaginaceae	49

0	<i>mmiphora habessinica</i> C	high cover value and characteristic species).
6	<i>ryaewellia</i> B	
0	<i>mucipouiteatroxylon</i> syM	
3	<i>hrenbergi</i> eansevieria S	
0	<i>illosav</i> Grewia	
0	<i>calypha fruticosa</i> A	
0	<i>eclea nobilis</i> T	
0	<i>aphia abyssinica</i> B	
2	<i>sucsembaz</i> Croton	
0	<i>uphorbia tirucalli</i> E	
2	<i>acia nilotica</i> A	
1	<i>imperischdgofera</i> nI	
0	<i>inifolius</i> xarfeenodiscus L	
8	<i>rua triphyllaea</i> M	
0	<i>abyssinica</i> ospyros D	
0	<i>capensis</i> Ol	
9	<i>lutinaev</i> Grewia	
8	<i>amrindus indica</i> T	
0	<i>donaea angustifolia</i> oD	
0	<i>rubi folius</i> Allophylus	
0	<i>ivorum</i> dea lcuE	
0	<i>arissa</i> Spinarum C	
0	<i>cokanthera schimperi</i> A	
0	<i>ium guineense</i> ygyz S	
0	<i>uropaeaea</i> Ol	
0	<i>calypha rasemosa</i> A	
0	<i>Slacia congolensis</i>	

Community and Species Diversity Indices

The Shannon diversity index (H) was a commonly used index to characterize species diversity in a community. Among the six communities, the Shannon – Wiener diversity index result shows community I to be the richest in number of species, whereas, community VI has the lowest number of species. Community I has the highest species diversity (2.798) with the highest number of species (144) followed by II, IV, V, VI and III with species diversity 2.381 (111), 2.360 (72), 1.606 (620), and 1.578 (71) respectively. Generally, diversity follows the trends observed in species richness (Whittaker, 1975). It is highest in species rich and lowest in species poor community. It has been proposed that the more numerous the species in a community, the greater the stability. In other words, high environmental stability leads to high community stability, which in turn permits high diversity of species (Bormann and Kellert, 1991). Accordingly, community types I, III and IV are in high stable environment and therefore, have high species richness. The high species richness is probably attributed to the optimum environment that supports the woodland species or attributed to the minimum level of disturbances. Species evenness measures the equality of species in a given sample area, or indicates lack of dominance by few species. It shows the relative proportional abundance of a species in communities. Low evenness value indicates the dominance of few species in the sample area. The community, which is relatively lower in species richness, appears to have high evenness. Based on this, communities III, VI and V have the lower evenness values. Community I has the highest evenness value. In this regard, communities I, IV and II have more even representation of all species. Jaccard's Coefficient of similarity is computed to see the similarity and dissimilarity of the six communities identified in the study area. Results of this analysis indicate that species composition similarity among the six communities is high. Communities I and II, V and VI, and II and III have relatively high similarity of species composition, whereas, I and V and I and VI have high dissimilarity of species composition. The most probably reason of species composition similarity or dissimilarity is the soil particles

This community exists at altitudes between 1188 m asl. *lypha fruticosa*, *Teclea nobilis*, *Baphis byssinica* Aca--1397 masl. *Croton zambescus*, *Euphorbia tirucalli*, *Acacia nilotica*, *Indigofera* are *Acacia tortilis* and *schimperi*, *Lecaniodiscus fraxinifolius*, dominant tree – shrub species. Less abundant but important species *Maerua triphylla*, *Diospyros abyssinica*, *Olea* in this community are *Grewia velutina*. and *capensis*, *subsp. macrocarpa* *Abrus* hand climber species in this community are also contain *precatorius*, *Glycine wightii*, *Monanthotaxis parvifolia*, *Cissus* with *Salacia congolensis* and *petiolata*, *Cynanchum altiscandens*, *Persitrophe paniculata*, lesser cover values. Species such as *Bidens* and *Gomphocarpus purpurascens*, *Momordica friesiorum*, are with relatively less surface cover. *pilosa*,

Maerua triphylla – *Diospyros abyssinica* – *Olea* Community V. *capensis* type.

Altitudinal distribution of this community type is found between 1280 masl– 1213 masl. Dominant tree species of this community . *1280 masl– 1213* *Baphia abyssinica*, *Lecaniodiscus fraxinifolius*, *Diospyros* were *Olea capensis*, *Maerua triphylla*, *Grewia velutina*, and *abyssinica*, *Dodonaea angustifolia*, *Euclea divinorum*, *Carissa spinarum*, *Tamarindus indica*, are tree–shrub species. *Acokanthera schimperi* and characteristic species of *Euclea ddivinorum* and *Carissa spynarum*, *Syzygium giuneense*, *Olea* this community. Species such as were with relatively high cover *Flacourtia indica* and *europaea*, *Ageratum conyzoides*, *Cynodon dactylon*, *Xanthium spinosum*, value. *Triumfetta rhomboida*, *Amaranthus caudatus*, *Centella asiatica*, are herbaceous surface layer species. *Lactuca inermis*

Croton zambescus – *Harrisonia abyssinica* – Community VI *Diospyros abyssinica* type.

This community type was distributed at altitudes between 1225– 1258 masl. is the characteristic shrub species of *Salacia congolensis* 1258 masl. *acalypha racemosa*, *Syzygium* the community. Species such as tree– shrub species *Flacourtia indica* and *guineense*, *Olea europaea*, with high cover value. On the other hand the following *Portulaca oleracea*, *Pupalia lappacea* *Ruellia prostrata*, species *Annonia senegalensis*, *Datura stramonium*, *Chenopodium schimperianum*, *Ficus vasta*, *Galinsoga parviflora*, *Hippocratea africana*, *Hygrophila schulli*, *Mollugo nudicaulis*, *Oncoba spinosa*, are only re-ordered at this *Ozoroa insignis* and *Ocimum urticifolium*, community in the entire study area.

IV 11	III 5	II 17	I 11	Community type cluster size
0.0	0.0	0.0	1.8	<i>anthium setiflorum</i> C
0.0	0.2	0.0	1.3	<i>cacia hockii</i> A
0.0	0.3	0.2	1.3	<i>appea capensis</i> P
2.0	0.6	0.9	1.4	<i>cacia brevispica</i> A
1.5	0.6	1.0	1.3	<i>bicolor</i> Grewia
1.7	1.3	0.2	1.9	<i>mbretum molleo</i> C
0.0	0.2	4.1	0.5	<i>mmiphora</i> Africanao C
0.5	1.6	4.0	1.3	<i>Rhus natalensis</i>
0.0	0.0	2.6	0.5	<i>cacia</i> Senegal A
0.1	0.7	2.0	1.0	<i>abyssinica</i> arrisonia H
0.0	0.0	1.2	0.2	<i>alanites aegyptiaca</i> B
0.0	0.0	1.1	0.6	<i>imenia Americana</i> X
0.0	0.4	1.5	0.3	<i>Ziziphus mucronata</i>
0.7	0.4	1.4	0.0	<i>ineriacicrostachys</i> D
0.8	3.4	0.2	0.5	<i>rminalia browniee</i> T
0.0	1.8	0.0	0.3	<i>anthozylon chalybeum</i> Z

able 2. Synoptic table of the Abaya – Hamassa vegetation with T diagnostic species having high cover value and characteristic species. (The figures printed in bold indicates species having

11-Discussion

Plant growth is influenced by physical and chemical properties soil, conducted environmental conditions and availability of water. These soil properties are important factors in determining plant growth and vegetation distribution. The test for variance made in present investigation for physical and chemical soil properties revealed that, the differentiation of communities can be partly explained by the variations of soil texture and physical and chemical parameters. The environmental data collected from Abaya-Hamassa were average in accordance with the community types identified and tested statistically for significant differences between communities. There was significant variation in particle size and distribution of sand, where as the variation in the size and percentage distribution of clay and silt particles was not so significant varying only 0.05 level of significance. Soil particle size distribution has a considerable effect on the vegetation of the study area affecting soil aeration, water movement, root penetration and water holding capacity. The top soil samples collected from 55 quadrats were categorized into 3 textural classes, sandy clay loam (Communities I and II), Sandy silt loam (communities III and IV), and sandy loam (communities V and VI). Soil texture was an important soil parameter that affects site quality. It influenced the nutrient supplying ability of soil solids, soil moisture, root development and air relations. Comparison of the community types based on the sand content of the soil had shown that the community type III is significantly different from communities II, IV, and V and is significantly different from communities I, II, IV, V and VI in its clay composition. Community type IV is significantly different from communities I, II, III, V and VI in its silt composition. Soil particles distribution along the communities was more or less similar. The proportion of sand in community III is the highest (52.2%) and the lowest (48.5%) in community V. The proportion of clay was high (27.09%) in community II and it was low (21%) in community III. The proportion silt was high (27.2%) in community IV and it was low (23.6%) in community I. In general, the variation of soil particle distribution among communities was small. It can be mooted that the variation of sand among communities was lowest when compared with clay and silt variation. Soil PH affects the growth of plants and the allocation of plant types by its effect on the availability of mineral nutrients and decomposition of organic matter. The average PH values of the six communities ranged between 6.58 – 7.12 values. Soils of communities II, III, IV, V and VI have neutral PH values whereas, community I is slightly basic. The neutrality and slightly basic character of the soil is due to low breakdown of organic matter and low leaching of soil attributable to the very low annual rainfall. There is no significant differences in soil PH except the slight variations obtained among the communities. The total nitrogen content of the soil was significantly varied in six communities. Soil of community V has higher total nitrogen content but community III had lower total nitrogen content. The mean values of total nitrogen in decreasing order of magnitude include communities V, IV, II, VI, I and III. In the total nitrogen content of the soil community type V was significantly different from community types III, I, VI, II and IV (Table-7). The total nitrogen content of the soil was determined by the amount of organic matter. It is higher in those community types with higher organic matter content and lowers in those having lower organic matter content. Climatic conditions especially temperature and rainfall, exerted a dominant influence on the amounts of nitrogen and organic matter in the soils. In general, the decomposition of organic matter was accelerated in warm climates. Dry soils are generally, low in the organic matter hence the total nitrogen content of the study area was low, since the area falls in between arid or semi arid region. The electrical conductivity of the soil solution was mainly determined by soluble salts of carbonates, bicarbonates, sulphates, chlorides and nitrates (Chopra and Kanwar, 1982). Electrical conductivity in communities II and III is the highest (3.60 and 3.62 mmhos/cm) respectively. It is lowest in community V (2.83

composition and the soil texture. The soil textural composition of communities I, II and III is sandy clay loam and that of communities V and VI sandy loam. The lowest species composition similarity is recorded between communities I and VI, that is 0.132 (13.2%). There is transition between communities I and VI and I and V. Comparatively small proportion of species occur in the two groups of communities in common (13.2%) and (16.8%) respectively. Generally, in all other communities the transition is gradual and species composition diversity is similar.

10-Similarity between vegetation of the study area with other vegetation types in Ethiopia.

The vegetation of Abaya Hamassa woodland can be compared with other vegetation types in Ethiopia. The vegetation of Key Afer Shala Luqa and Southwest of Lake Chamo (Teshome Soromessa, 1997), Vegetation of Lake Abaya to Chench Highland (Desalegn Wana, 2002), Rangeland Vegetations of Alagae and Neteli (Ali Seid, 2004), Vegetation of dry land part of North Shewa (Hussien Adal, 2004), Vegetation of the Savanna grassland and woodland in Nechisar National Park (Tamrat Andargie, 2001) and Vegetation of semi-wetland of Cheffa area South Wello (Bayafer Tamene, 2000) differ significantly with some sort of similarities with the vegetation types in the present study (Table-8). Key Afer-Shala Luqa Vegetation is located about 709 km south-west of Addis Ababa and 200 km South of Arbaminch between longitudes 370 20' and 370 32' E and latitudes 50 41' and 50 56' N. Key Afer-Shala Luqa lies between 600-1900m.a.s.l. in altitude. Southwest of lake Chamo is found between longitudes 360 34' and 360 56' E and latitudes 50 15' and 50 40' N which is about 530 km from Addis Ababa. Altitudinal ranges between 1100m.a.s.l. around Lake Chamo and 1900m.a.s.l. around Zeyise. The vegetation of Lake Abaya to Chench highlands is located 60 05' and 60 12' N and longitudes 370 33' and 370 39' E. Altitudinal ranges between 1177m.a.s.l. around Lake Abaya and 2718m.a.s.l. in Chench highlands. Rangeland Vegetations of Alagae and Neteli is located within the Ziway-Shalla drainage basin in latitude 70 00' and 70 30' N and longitudes 380 00' and 380 30' E. the altitude of the area ranges from 1540m.a.s.l. to 2075m.a.s.l. The vegetation along the Eastern escarpment of Wello has an altitudinal range of 1660-1900m.a.s.l. Dry land vegetation of North Shewa is situated between latitudes 80 38' and 100 42' N and longitudes 380 40' and 400 30' E. The elevation of the area ranges between 1050-1290m.a.s.l. Nachisar National park is located near Arbaminch town, 505 km South of Addis Ababa. It is positioned at the center of the Ethiopian Rift Valley between latitudes 50 55' and 60 05' N and longitudes 370 48' E of the equator. The comparison was based on the similarities in distribution. A similarity analysis was carried out based on the presence of species in order to evaluate the relationship between these vegetations. The similarity index used was Sørensen's Similarity Coefficient $2c/a+b$ where c is the number of species shared by the vegetations compared, a is the number of species in one vegetation and b is the number of species in the other vegetation. The vegetation in Abaya-Hamassa was floristically related more to the vegetation of key Afer-Shala Luqa and Southwest of Lake Chamo, Vegetation along the Eastern Escarpment of Wello and Vegetation of Semi-Wetland of Cheffa area South Wello than to the Rangeland Vegetation of Alagae and Neteli and Dry land Vegetation of North Shewa. The species composition of Vegetation of Nechisar National Park and Vegetation of Lake Abaya to Chench Highlands were relatively similar to vegetation of Abaya-Hamassa. However, Range land vegetation of Alagae and Neteli and Dry land Vegetation of North Shewa show some variation may be due to vegetation disturbance. Species those which were common in all these areas *Aca-cia seyal*, *A. tortillis*, *Acalypha fruticosa*, *Achyranthes* were: *asperra*, *Acokanthera schimperi*, *Asparagus africanus*, *Balanites aegyptiaca*, *Barteria eranthemoides*, *Cadaba farinose*, *Cissus spp.*, *Combretum spp.*, *commiphora spp.*, *Ficus spp.*, *Grewia spp.*, *Cyperus spp.*, *Rhus natalensis*, *Ximenia americana* and *Ziziphus spp.*

factors such as moisture, ambient temperature, disturbance and dispersal effects in community organization and diversity.

Table 3. Shannon – Wiener Diversity index calculated among the 6 communities.

Evenness (H')	Diversity Index (H)	Richness	Community
0.563	2.798	144	I
0.509	2.381	111	II
0.370	1.578	71	III
0.501	2.360	107	IV
0.415	1.775	72	V
0.389	1.606	62	VI

Table 4. A presence half matrix of Jaccard's coefficients calculated among 6 communities.

VI	V	IV	III	II	I	Community
					1.00	I
				1.00	0.28	II
			1.00	0.27	0.24	III
		1.00	0.24	0.24	0.21	IV
	1.00	0.21	0.22	0.21	0.16	V
1.00	0.27	0.22	0.21	0.22	0.13	VI

Table 5. Jaccard's community coefficient among adjacent communities/altitudinal ranges

Communities	Similarity coefficient	Altitudinal ranges
I, VI	13.2%	1241 - 1410/1225 – 1258
I, V	16.8%	1214 - 1410/1213 – 1280
I, III	27.7%	1184 – 1309/1233 – 1283
V, VI	27.9%	1213 – 1280/1225 – 1258
I, II	28.1%	1241 – 1410/1184 - 1309

Table 6. Average values of environmental variables (indicate

Texture	tlis %	Clay %	ndaS %	TN	E (C)	HP	eopIS (0)	Ita eduit m)(SD)
Sandy clay loam	23.59	26.41	50.00	0.54	3.28	7.12	6.65	1242.42	I
Sandy clay loam	7.47	14.23	17.47	0.48	1.09	0.48	8.02	30.89	
Sandy clay loam	24.45	27.09	48.45	0.84	3.62	6.67	12.09	1288.18	II
Sandy clay loam	7.24	9.60	13.43	0.89	0.97	0.53	10.60	49.75	
Sandy silt loam	26.80	21.00	52.20	0.36	3.60	6.58	15.20	1257.80	III
Sandy silt loam	6.53	5.66	9.91	0.24	0.24	0.32	11.95	22.02	
Sandy silt	27.22	24.11	48.67	1.02	3.13	6.83	9.89	1244.78	IV

mmhos/cm).Vegetation reacts variously to exposure; light loving communities (Heliophytes) reach their highest limits on southern exposure, while shade plants, shade loving communities reach their high- est limits on north, northeast and northwest exposures. Light absorption between south and north slopes in the mountain is very considerable, much greater than in the plains. The total light in a south exposure reaches 1.6 to 2.3 times higher values than in a north exposure. The effect of slope aspect as a control of radiation may not appear to be pronounced due to the low variation of altitudinal gradients among communities. Therefore, it has no significance in determining the distribution of the vegetation.The slope of the soil surface affects vegetation directly as well as indirectly. The indirect effect was due to the influence up on the water supply of the soil and the shifting of the angle of incidence of the sun's rays, modifying the intensity of insulation. The effect of slope was seen in its influence on the run-off and drainage and consequently upon the nutrient, depth and water content of the soil. In the present study slope had shown a significant difference in the distribution of the communities. Community type I had gentle slope with high species diversity. This could partly be due to high retention of water and moderate composition of soil particles. Community type III had steep slope, and fairly high percentage surface stone cover with low species diversity. This may be due to less retention of water in the soil for having steep slope. Community types II and V have same mean values of slope. But both communities had different species diversity.The diversity index calculated for the 6 communities had shown that the species diversity of community type II was 2.381 and the species diversity of community type V was 1.775.The difference in species diversity in community type II and in community type V may occurred as of different amounts of total nitrogen content in the soils of two communities. The mean values of total nitrogen content of community type II was 0.84 and that of community type V was 1.72. High species diversity in community type II could partly be due to the low available total nitrogen content in the soil. This agrees with the generally held belief that highly nutrient rich soils would tend to support low species diversity (Huston, 1994). This is the reason that community type II supports high species diversity with low nutrient content in the soil, while community type V supports low species diversity with high nutrient content in the soil. Since both communities have the same averaged values of slope.Altitude is an important environmental factor that affects atmospheric pressure, moisture, and temperature, and temperature which has a strong influence on the growth of plants and the distribution of vegetation. The mean altitudinal variation among 6 communities identified and was significant at $P < 0.05$ level of significance. The mean values of altitude in communities range between 1239.60 to 1288.18m.a.s.l.In order to examine the significant differences and/or similarities between the community types identified, Pearson's product moment correlation coefficient comparison test was employed for the environmental parameters. There was significant correlation between percentage (%) with total nitrogen. All other environmental parameters were correlated insignificantly at $P < 0.05$ level of significance. Altitude was negatively correlated with % of sand and % of clay. Except these correlations, all the other environmental variables were positively correlated. With increasing altitude, the steepness of the surface increases and this results in the loss of soil particles. This may be the reason that altitude was negatively correlated with percentage of sand and clay.Based on the findings, it was suggested that the present trend of heavy exploitation of plant species in the area reducing the vegetation cover and may lead to local extinction of species. Hence a proper exercise of conservation practice that which can promote natural regeneration based on well worked out management plans should be adopted at the earliest possible time. The conservation of this vegetation types should be given the priority concern for the conservation of biodiversity. It was also suggested that further investigation of the vegetation–environment relationships is essential by considering

Sandy loam	25.17	25.33	48.50	1.72	2.83	6.75	12.17	1241.33	V	
	5.42	12.16	14.68	1.14	0.44	0.23	5.12	26.93		
Sandy loam	24.40	25.00	50.60	0.76	3.30	6.70	9.80	1239.60	VI	
	9.34	11.79	19.37	0.74	0.72	0.20	9.96	12.93		

loan 5.88.00.2281.00.50.49
37.33

Table 7. Pearson's product moment correlation coefficient for correlations (bold values are significant at $\alpha = 0.05$). NS = not significant; S = significant; EC = electric conductivity; TN= total nitrogen.

Silt	Clay	Sand	TN	EC	PH	Slope	Altitude	
							1	Altitude
						0.235		
					1	Ns		Slope
						0.653		
					0.786	0.253		
					1	Ns	Ns	PH
						0.064	0.628	
					0.632	0.314	0.049	
				1	Ns	Ns	Ns	EC
					0.178	0.544	0.926	
				0.753	0.253	0.038	0.017	
			1	Ns	Ns	Ns	Ns	TN
				0.084	0.629	0.943	0.975	
			0.848	0.510	0.425	0.276	-0.273	Sand
		1	Ns	Ns	Ns	Ns	Ns	
			0.033	0.302	0.400	0.597	0.600	
		0.738	0.446	0.235	0.471	0.641	-0.039	Clay
	1	Ns	Ns	Ns	Ns	Ns	Ns	
		0.094	0.375	0.654	0.346	0.171	0.942	
	0.805	0.231	0.029	0.084	0.375	0.654	0.291	
1	Ns	Ns	Ns	Ns	Ns	Ns	Ns	Silt
	0.053	0.659	0.957	0.875	0.464	0.159	0.576	

Table 8. Similarities of vegetation between Abaya and other natural vegetation in Ethiopia.

Here, N = number of species included in comparison, C = number of species in common, S = Sørensen's coefficient of similarity.

S	C	N	Vegetation
0.54	93	216	Vegetation of Key Afer-Shala Luqua and

- Edwards, S. (1992). Some threatened endemic plants of Ethiopia In: (Sue Edwards and Zemede Asfaw eds.), Botany 2000: East and Central Africa. NAPRECA Monograph Series No. 2., pp. 35-55.
- (1980). Assistance to land use, Planning experts' final report. **FAO**. 8 Forest resources and potentials for development (Ethiopia). AGOA:Eth/78/003, UNDP/FAO, Addis Ababa. PP. 54.
- (1990). Soil Map of the World: Revised Legend. **FAO/UNESCO**, 9 FAO/UNESCO, Rome.
- (1983). Quantitative plant ecology. Third ed. **Greig-Smith, P.** 10 Butterworths, London.
- (1992). Global biodiversity status of the earth's **Groombridge, B.** 11 living resources. A report compiled by the world Conservation Monitoring Center Chapman and Hall. London.
- (2004). Traditional use, management and conservation **Hussien Adal**. 12 of useful plants in dry land parts of North Shewa zone. Msc. Thesis unpublished. Addis Ababa University Huston, M. A. (1994). Biological Diversity. The coexistence of species on changing landscapes. Cambridge University press. Cambridge.
- (1973). Soil chemical analysis, Prentice Hall of India **Jackson, M. L.** 13 Private limited New Delhi.
- (1978). Selected methods for soil and plant analysis. Institute **Juo, A.** 14 of Tropical Agriculture. Ibadan, Nigeria.
- (1973). Quantitative and dynamic plant **Kershaw, K. A.** 15 ecology. 2nd ed. The English Book Society and Edward Arnold publishers Ltd. London.
- (1981). Modern concept of ecology. 2nd (nd) reviewed **Kumer, H. D.** 16 ed. Vikas publisher, Delhi.
- (1971). The Geography of Ethiopia. University college of **Mohr, P. A.** 17 - Addis Ababa. Central printing press. Addis Ababa, Ethiopia. Mueller
- Dombois, D. and Ellenberg, H. (1974). Aims and Methods of Vegetation Ecology.
- (1993). Ecology 2. John **John Wiles and Sons Inc.USA Paul, C.** 18 Wiley and Sons Inc. New York Brisbane. Toronto. Singapore.
- (1949). The mathematical theory of **Shannon, C. E. and Wiener, W.** 19 communication. University of Illinois press. Urbana III.
- (1971). The description and classification of **Shimwell, D. W.** 20 vegetation. Sidgwick and Jacson. London.
- (1997). An ecological study of the lowland **Teshome Soromessa**. 21 vegetation of Key Afer-Shala Luka and southwest of Lake Chamo. Msc. thesis unpublished. Addis Ababa University.
- (2001). Floristic composition and Ecology of the **Tamrat Andargie**. 22 Savanna grassland and Woodland vegetation in Nechisar National Park and its conservation status. Msc. thesis unpublished. Addis Ababa University. U. S.
- (1992). Conserving Biodiversity. A **National Research Council**. 23 research agenda for Development Agencies.
- Washington D. C. Vaukasimovic, S. (1969). **National academy Press**, 24 Report and Interpretation of the pedological investigation in the Harre Region, Near Arbaminch Gamo Gofa Province, Zurich (Switzerland).
- (1979). Multivariate methods in phytosociology **van der Maarel, E.** 25 with reference to the Ntherland. The study of vegetation. Junk.
- (1991). Biodiversity Guide to Ethiopia. World **The Hague. WCMC**, 26 Conservation Monitoring Center. Whittaker, R. H. (1962). Classification of plant communities. Biological Rev. 28: 1-239.
- (1983). The vegetation of Africa: A description Memoir to **White, F.** 27 accompany the UNESCO/AETFAT/UNESCO vegetation map of Africa. Natural Resource Research 20. UNESCO. Paris.
- (1967). Gradient analysis of vegetation. Biol. Rev. 1: **Whittaker, R. H.** 28 281-289.
- (2013). Annual report to the **Wolayta zone Agricultural office**. 29 Southern Nations, Nationalities and Peoples' Regional State Agricultural Office.
- and Mesfin Tadesse (1990). The status of the **Zerihun Woldu**. 30 vegetation in the lake regions of the rift valley of Ethiopia and the possibilities of its recovery. SINET: Ethio. J. Sci. 13: 97-120.
- and Asferachew Abate (1999). **Zerihun Woldu, Mesresha fetene**. 31 Vegetation under different tree species in Acacia woodland in the rift valley of Ethiopia. SINET: Ethio. J. Sci. 22: 235-252 Faculty of Science Addis Ababa university.

0.43	Southeast of Lake Chamo		Vegetation of Lake Abaya to Chench highland
0.33	65	213	Rangeland Vegetation of Alagae and Neteli
0.58	97	216	Vegetation along the eastern escarpment of Wello
0.34	67	208	Dryland vegetation of North Shoa.
0.48	83	199	The Savana grassland and woodland Vegetations in Nechisar National Park
0.50	87	206	Vegetation of semi-wetland of cheffa area South Wello

استنتاج

Based on the findings, it was suggested that the present trend of heavy exploitation of plant species in the area reducing the vegetation cover and may lead to local extinction of species. Hence a proper exercise of conservation practice that which can promote natural regeneration based on well worked out management plans should be adopted at the earliest possible time. The conservation of this vegetation types should be given the priority concern for the conservation of biodiversity. It was also suggested that further investigation of the vegetation-environment relationships is essential by considering factors such as moisture, ambient temperature, disturbance and dispersal effects in community organization and diversity.

ref_str

- (2004). Vegetation description and condition assessment of **li Seida**. 1 protected rangelands of Alagae and Neteli. Msc. thesis unpublished. Addis Ababa University.
- (2000). A floristic analysis and Ethnobotanical **Bayafers Tamene**. 2 study of the semi-wetland of chefa area South Wello. Msc. thesis unpublished. Addis Ababa University.
- and Kellert, S. R. (ed.) (1991). Ecology, Economics, **Bormann, F. H.**, 3 Ethics. The broken circle. Yale University press New Haven and London.
- and Kanwar, J. S. (1982). Analytical Agricultural **Chopra, S. L.** 4 Chemistry. Kalyani publishers, New Delhi.
- (1994). The Borana plateau of Southern Etiopia: **Coppock, D. L.** 5 Synthesis of pastoral Research, Development and Change, 1980-1991. ILCA. Addis Ababa, Ethiopia.
- (2002). Plant communities and diversity along **Desalegn Wana**. 6 altitudinal gradients From Lake Abaya to Chench highlands. MA. Thesis unpublished. Addis Ababa University.
- Zerihun Woldu and **Ensermu Kelbessa, Sebsebe Demissew**, 7



IJSURP Publishing Academy

International Journal Of Scientific And University Research Publication
Multi-Subject Journal

Editor.

International Journal Of Scientific And University Research Publication



+965 99549511



+90 5374545296



+961 03236496



+44 (0)203 197 6676

www.ijsurp.com