



International Journal Of Scientific And University Research Publication

ISSN No **2364/2018**

Listed & Index with
ISSN Directory, Paris



Multi-Subject Journal



EFFECT OF NICKEL ON ANTIOXIDANT ENZYME (SUPEROXIDE DISMUTASE AND CATALASE) OF *KIDDERYUS MELANOSTICTUS*

Puspanjali Parida || P. G. Department of Zoology
North Orissa University
Baripada
Odisha
757003 India

Duttaphrynus melanostictus were divided into four groups as A, B, C and D. Each group consisting of 10 animals. Animals of group A (control) were given orally 50 µl of

distilled water. The experimental animals of group B, C and D were administered orally with 50 µl (=50 µg) of nickel. The animals were sacrificed after different time intervals such as 0h (group A), 24h (group B), 48h (group C) and 72h (group D). The superoxide dismutase and catalase of kidney at different time intervals were measured and compared.

Duttaphrynus melanostictus, Nickel,
 kidney, superoxide dismutase and

intervals. The animals were sacrificed at 24 hours, 48 hours and 72 hours and different parameters were measured.

مقدمة

Preparation of tissue sample

(both control and *Duttaphrynus melanostictus* Body weight of experimental) was measured by digital monopan balance (Shimadzu; ELB 300) and were sacrificed at 0 hr, 24 hr, 48hr and 72 hr of time interval. The kidney was dissected out quickly and kept at 0°C. A 20% homogenate was prepared in ice-cold 50 mM phosphate buffer (pH 7.4) using pre-chilled porcelain mortar and pestle by up and down strokes at 40°C. The homogenate was centrifuged at 4000 rpm (1000Xg) for 10 minutes at 40°C in Cooling Centrifuge (Remi). The supernatant was taken for biochemical assay.

Measurement of protein content

Protein estimation of samples was made according to the method of Lowry (1961). The data were expressed in mg/g tissue.

Measurement of superoxide dismutase (SOD) activity

Superoxide dismutase (SOD; EC 1.15.1.1) activity was determined (2000). SOD activity was expressed as units/mg protein.

Measurement of catalase (CAT) activity

Catalase (CAT; EC 1.11.1.6) activity was estimated according to Beers and Sizer (1952). The activity of catalase was expressed as nkat/mg protein (1 nkat=1 mole of substrate converted to product per sec, 1U=16.67 nkat).

Statistical methods

One way ANOVA and Post Hoc analysis was carried out to find out treated *Duttaphrynus melanostictus* the level of significance between with nickel over a period of 24 hr, 48 hr, 72 hr and in control. A difference was taken as significant when P was less than 0.05. Statistics is done with the help of software SPSS pack age 16.0.

Body weight and colour of kidney

exposed to nickel were *Duttaphrynus melanostictus* Body weight of 96.6±14.33 before dose (BD) and 94.6±14.33 after dose (AD) in 24 hour, 95.4±16.42 before dose and 92.2±16.08 after dose in 48 hour, 95.6 ± 27.46 before dose and 90.6 ± 27.46 after dose in 72 hour. In after dose *Duttaphrynus melanostictus* other words body weight of decreased maximum in 72 hour (Fig.5). The colour of kidney varies at different time intervals (Figs 1, 2, 3,4).

Protein content

treated *Duttaphrynus melanostictus* Protein content (mg/g tissue) in 1.80 mg/g tissue ± 1.59 mg/g tissue, 22.03± with nickel were 22.33 0.90 mg/g tissue IJSR ± 1.24 mg/g tissue, 13.28±, 28.60

of contaminants are continuously introduced into the aquatic environment mainly due to increased industrialization, technological development, growing human population, oil -exploration and exploitation, agricultural and domestic wastes run (Lima constitute one of the most dangerous groups because of their persistent nature, toxicity, tendency to accumulate in organisms and undergo food chain amplification and more still, they are nondegradable (Fufeyin and Egborge, 1998). Antioxidant defense enzymes (ADS) play a crucial role in maintaining cell homeostasis. ADS may be induced after exposure to pollutants, this response reflecting an adaptation of the species to their environment. This system may also be inhibited, which may lead to antioxidant mediated toxicities. Some heavy metals are hepatotoxic agents causing liver disorders, largely due to their active metabolites and free radicals. In small quantities, certain heavy metals are nutritionally essential for a healthy life. Some of these are trace elements (e.g., iron, copper, manganese, and zinc). These elements, or some form of them, are commonly found naturally in foodstuffs, fruits and vegetables, and in commercially available multivitamin products. Diagnostic medical applications include direct injection of gallium during radiological procedures, dosing with chromium in parenteral nutrition mixtures, and the use of lead as a radiation shield around xray equipment (Roberts, 1999). Animal studies have reported reproductive and developmental effects, such as a decreased number of live pups per litter, increased pup mortality, and reduction in fetal body weight, and effects to the dam from oral exposure to soluble salts of nickel (U.S. EPA IRIS 1999) In the present work generation of ROS in response to nickel were estimated by measuring Protein content, Superoxide dismutase (SOD) and Catalase (CAT) at different time *Duttaphrynus melanostictus* activity in kidney of intervals (24 hr, 48hr and 72hr) and compared with the control.

Duttaphrynus melanostictus (AS DHOT MATERIALS AND METHODS)

(80 g to 110) g were collected during *uttaphrynus melanostictus* D night and early morning time locally in Baripada from August 2014 to November 2014. They were acclimatized for seven days prior to the experiment. The animals were kept in plastic perforated jar for experiment. The animals were divided in to two groups, i.e., control and experimental.

Preparation of stock solution

The stock solution was prepared by dissolving 1mg of nickel chloride in 1ml of distilled water. From this stock solution 50µl (50µg 50µl of .nickel chloride) was taken and given orally to the animal nickel were given orally to 15 numbers of toads at different time

Volume : 5 | Issue : 5 | May 2016 • ISSN No 2277 - 8179 | IF : 3.508 | IC Value : 69.48 highest at 24 hours (Fig. 8). One way ANOVA *Duttaphrynus* revealed that the CAT activity (nkat/mg protein) in exposed to nickel at different time intervals is *melanostictus* significant [F(3, 19) = .665, P = .585]. Post Hoc analysis revealed that CAT activity (nkat/mg protein) at different time intervals when was significant at *Duttaphrynus melanostictus* treated with nickel in 24 hour and 48 hours (P < 0.05; LSD). However, not significant at 72 hour with respect to the control.

NOISSISCUD

It has been demonstrated that the organophosphates diazinon and chlorpyrifos both reduce rates of metamorphosis in tadpoles of the *et* (Sumanadasa *Duttaphrynus melanostictus* Asian common toad *Duttaphrynus* 2008). In our study we found the body weight of *al.*, exposed to nickel significantly decreased than that of *melanostictus* controls. At 24 hour, 48 hour and 72 hours time intervals of nickel exposure, the body weight of toads were decreasing with increase of time and also change in the colour of kidney. The two important antioxidant enzyme SOD and CAT increases at 24 hour indicate efficient mechanism of kidney in response to oxidative stress generated in response to Nickel Chloride. At 72 h the animal almost neutralize the level of SOD and CAT.

استنتاج

From this work it is known that the animal has well equipped with two important antioxidant enzyme SOD and CAT to neutralize the environmental oxidative stress for its survival.

ref_str

- Lima D., Santos M. M., Ferreira, A. M., Micaelo C. and .1
Reis Henriques., (2008) The use of the shanny *Lipophrys pholis* for
pollution monitoring; A new sentinel species for the northwestern
European marine ecosystems. *Environment International*, 34: 94 –
101.
- Fufeyin, T. P. and Egborge A. B. M., (1998) Heavy metals of Ikpoba .2
River, Benue State, Nigeria. *Tropical Freshwater Biology*, 7: 27 – 36.
- Training Manual on Roberts, J. R., (1999) Metal toxicity in children. In: .3
Emeryville, *Pediatric Environmental Health: Putting It into Practice*
CA: Children's Environmental Health
pdf/ ([http://www.cehn.org/cehn/trainingmanual/Network
manualfull.pdf](http://www.cehn.org/cehn/trainingmanual/Networkmanualfull.pdf)).
- Integrated Risk U.S. Environmental Protection Agency., (1999).4
National Center for Information System (IRIS) on Nickel Refinery Dust
Environmental Assessment, Office of Research and Development,
Washington, DC.
- Das, K., Samanta, L. and Chainy, G. B. N., (2000) A modified .5
spectrophotometric assay of superoxide dismutase using nitrite
, 37: *Ind. J. Biochem. Biophysics* formation by superoxide radicals.
201-204.
- Beers, R. F., Jr. and Sizer, I. W., (1952) A spectrophotometric .6
method for measuring the breakdown of hydrogen peroxide by
, 195: 133-140. *J. Biol. Chem.* catalase.
- Lowry, O. H., Rosebrough, N. J., Farr, A. L. and Randall, R. J., .7
J. Biol. (1951) Protein measurement with the Folin phenol reagent.
, 193: 265-275. *Chem.*
- Sumanadasa D.M., Wijesinghe M.R. and Ratnasooriya W.D., (2008).8
Bufo Effects of Diazinon on larvae of the Asian Common Toad (
Environmental Toxicology and Schneider 1799). *melanostictus*
27 (11): 2320-2325. *Chemistry*

INTERNATIONAL JOURNAL OF SCIENTIFIC RESEARCH 409
Volume : 5 | Issue : 5 | May 2016 • ISSN No 2277 - 8179 | IF : 3.508 | IC Value : 69.48 after 0 hour, 24 hour, 48 hour and 72 hour respectively. Protein content (mg/g tissue) gradually decreased at 0 hour, 24 hour and 72 hours. It was lower in 72 hour in comparison exposed to nickel at different time *Duttaphrynus melanostictus* to intervals. The protein content was highest at 48 hours (Fig.6). One way ANOVA was performed in order to analyse the effect of nickel *Duttaphrynus* on the protein content at different time intervals in . One way ANOVA revealed that the protein content *melanostictus* is significant *Duttaphrynus melanostictus* at different time intervals in [F(3,18) = 135.212, P = .000]. Post Hoc analysis revealed that the protein content at different time intervals when treated with nickel was significant at 24 hour, 48 hour and *Duttaphrynus melanostictus* in 72 hour (P < 0.05; LSD).

Superoxide dismutase (SOD) activity

Duttaphrynus Superoxide dismutase activity (unit/mg protein) in 47.46 unit/mg \pm treated with nickel we were 34.09 *melanostictus* 124.55 unit/mg protein after 24 hours, \pm protein at 0 hour, 109.73 42.30 \pm 56.51 unit/mg protein after 48 hours, 38.47 \pm 51.31 unit/mg protein after 72 hours. The SOD level (unit/mg protein) *Duttaphrynus* lower in control in comparison to exposed to nickel at different time intervals. The SOD *melanostictus* level was highest at 24 hours and then gradually decreased at 48 hours and 72 hours (Fig. 7). One way ANOVA revealed that the SOD exposed to *Duttaphrynus melanostictus* activity (unit/mg protein) in nickel at different time intervals is significant [F(3, 19) = 1.076, P = .387]. Post Hoc analysis revealed that the SOD activity (Unit/mg protein) at different time intervals when treated with nickel was significant Only at 24 hour (P *Duttaphrynus melanostictus* in <0.05; LSD). While 48 hour and 72 hour are not significant with respect to control.

g 1. Untreated kidney of *Duttaphrynus melanostictus* iF

Fig 2. Kidney colour became more dark at 24 hour in nickel treated

Duttaphrynus melanostictus

Research Paper

g 3. Kidney colour became slightly reddish at 48 hour in nickel iF

uttaphrynus melanostictus D treated

g 4. Kidney colour normal at 72 hour in nickel treated iF

Duttaphrynus melanostictus

g.5: Comparison of body weight in *Duttaphrynus* iF *melanostictus* treated with Nickel at different time interval.

410 IJSR - INTERNATIONAL JOURNAL OF SCIENTIFIC RESEARCH

Research Paper

g.6: Comparison of Protein content activity (mg/g tissue) in iF *Duttaphrynus melanostictus* treated with Nickel at different time interval.

g.7: Comparison of SOD activity (unit/mg protein) in iF *Duttaphrynus melanostictus* treated with Nickel at different time interval.

g.8: Comparison of CAT activity (nkat/mg protein) in iF *Duttaphrynus melanostictus* treated with Nickel at different time interval.

Catalase (CAT) activity

Duttaphrynus Catalase activity (nkat/ mg protein) in 0.55 nkat/mg \pm treated with nickel were 4.02 *melanostictus* 0.59 nkat/ mg \pm 1.015 nkat/ mg protein, 4.42 \pm protein, 4.46 0.45 nkat/ mg protein after 0 hour, 24 hours, 48 hours \pm protein, 3.99 and 72 hours respectively. The CAT level (nkat/ mg protein) was lower in 0 hour in comparison to 24 hour and then gradually decreased at 48 hours and 72 hours. The CAT level (nkat/ mg exposed to nickel was *melanostictus* protein) of *Duttaphrynus*



IJSURP Publishing Academy

International Journal Of Scientific And University Research Publication
Multi-Subject Journal

Editor.

International Journal Of Scientific And University Research Publication



+965 99549511



+90 5374545296



+961 03236496



+44 (0)203 197 6676

www.ijsurp.com