



International Journal Of Scientific And University Research Publication

ISSN No **122/350**

Listed & Index with
ISSN Directory, Paris



Multi-Subject Journal

Volum : **(8)** | Issue : **(10)** |



CASE REPORT - ORAL LICHEN PLANUS IN A TEENAGER WITH CARDIOFACIOCUTANEOUS SYNDROME

Dr. Mariana Silvia Gandolfo
|| Associate Professor

Cardiofaciocutaneous syndrome (CFCS) is an extremely rare genetic disorder that belongs to syndromes known as RASopathies. The craniofacial and ectodermal manifestations of this syndrome are distinctive and play an important role in the diagnosis of this entity. Nevertheless, there is no scientific evidence to date of the presence of CFCS-related lesions in the oral mucosa. The present study is the first to report the presence of oral lichen planus (OLP) in a patient with CFCS. This finding should be taken into account in the assessment and diagnosis of CFCS in order to establish whether it is an un-related lesion or is associated with the syndrome.

Cardiofaciocutaneous syndrome, RAS genes, RAS proteins, RASopathies,

rough dry skin on the outer surface of her thighs (keratosis pilaris), tall forehead, bitemporal constriction, hypertelorism, and low-set ears. White patches and fissures were observed at both commissures of the lips (Figures 1 and 2).

مقدمة

Cardiofaciocutaneous syndrome (CFCS) is a genetically heterogeneous clinical entity that is inherited in a dominant autosomal manner. It belongs to a group of syndromes known as




Figure 1: A) Bitemporal constriction and low-set ears. Commissures of the lips: White patches and fissures B) Hypertelorism and very sparse eyebrows.



Figure 2: A) Dry rough skin B) Keratosis Pilaris C) Plantar aspect of the foot: plantar keratosis on areas subjected to pressure.

Examination of the oral cavity revealed the presence of white patches, plaques and verrucous lesions in the anterior third of the cheek mucosa on both sides. Red atrophic erythematous patches, alternating with white plaques, were observed in the attached and unattached gingiva of both jaws. A keratotic white plaque and the presence of a verrucous area were observed in the middle third of the dorsal aspect of the tongue. Ogival palate and marked tooth crowding were also observed (Figures 3, 4, 5).

(reference sequence NM_004333.4) in peripheral blood DNA showed the presence of variant c.1467>C (p.Leu489Phe) in exon 14 of gene RAF1 in heterozygous condition. The studied exons and flanking regions included the regions with the highest frequency of mutations. The detected variant has not been described previously. In silico prediction analyses indicated a potential deleterious effect of the detected variant on RAF1 function. Because the patient's mother also showed clinical features of the syndrome, the corresponding genetic sequencing was carried out, and showed the presence of variant c.1467>C (p.Leu489Phe) in exon 14 of gene RAF1 in heterozygous condition. The patient was the first offspring of a nonconsanguineous couple, and perinatal history and delivery were uneventful. The presence of typical cutaneous anomalies of the syndrome were diagnosed at the age of 3 years, and oral mucosa lesions were detected at the age of 10 years by her dentist, who referred her to our Service. Physical examination revealed dysmorphic facial features, sparse dry brittle hair, scant eyebrows,

Oral lichen planus (OLP) is a chronic inflammatory disease of immune nature and of unknown etiology usually presenting in middle age, and that can affect the skin, oral and genital mucosa, scalp, and nails. It has diverse clinical presentations, the most frequent being the variant observed in our patient: hyperkeratotic lichen (9). The prevalence of cutaneous lichen planus in the adult population worldwide ranges from 0.2% to 1%, whereas OLP is more frequent and has been reported to affect 1 to 4% of the population (9, 10). Overall, it affects women more often than men in a ratio of 1.5:1, and most cases develop between the age of 30 and 60 years (11, 12). In this regard, our patient was female and lichen planus occurred in the oral mucosa, both frequent observations and both in line with studies published in the literature. However, in contrast with reported findings that OLP is rare in children, with 5 to 10% of cases occurring in this age group (10, 13), the patient reported here was only 12 years old.

Figure 3. A) Tooth crowding B) Sectors I and II: White patches and plaques

Although CFCS is an extremely rare genetic disorder, it must be suspected in individuals presenting the phenotypical features described above. Craniofacial and ectodermal manifestations of this syndrome are distinct and play an important role in the diagnosis of this entity. Nevertheless, there is no scientific evidence to date of the presence of lesions in the oral mucosa associated with CFCS. In this regard, it would be important to perform thorough inspection of the oral mucosa of CFCS patients to record the presence or absence of stomatological lesions, in order to determine whether these lesions are another manifestation of CFCS. In addition, it is important to point out that OLP undergoes malignant transformation in 1 to 2% of cases, so that early diagnosis is of great clinical significance in terms of prognosis and patient follow-up (14, 15).



Figure 4: A) Upper gingiva: white and red patches B) Lower gingiva: white plaques C) Anterior third of the cheek mucosa: white plaque-like lesion and verrucous area D) Dorsal aspect of the tongue: keratotic white plaque and verrucous area.



Figure 5: A) Gingiva: white patches and plaques B) Ogival palate. Palatal gingiva: white patches and plaques

استنتاج

The present study is the first to report the presence of OLP in a patient with CFCS. This finding should be taken into account when evaluating and diagnosing CFCS, with the aim to establish whether it is an occasional manifestation or whether it is associated with the syndrome.

Conflict of interest •

The authors declare no conflicts of interest.

ref_str

[1] Tidyman WE, Rauen KA. Noonan, Costello and cardiofacio-cutaneous syndromes: dysregulation of the RasMAPK pathway. *Expert Rev Mol Med.* 2008; 10:e37.

[2] Tidyman WE, Rauen KA. The RASopathies: developmental syndromes of Ras/MAPK pathway dysregulation. *Curr Opin Genet Dev.* 2009; 19:230–6.

[3] Niihori T, Aoki Y, Narumi Y, et al. Germline KRAS and BRAF mutations in cardio-facio-cutaneous syndrome. *Nat Genet.* 2006; 38:294–6.

[4] Rodriguez-Viciano P, Tetsu O, Tidyman WE, et al. Germline mutations in genes within the MAPK pathway cause cardio-facio-cutaneous syndrome. *Science.* 2006; 311: 287–90.

[5] Roberts AE, Allanson J, Jadico SK, Kavamura MI, Noonan J, Opitz JM, et al. The cardiofaciocutaneous syndrome. *J Med Genet.* 2006;43:833–42.

[6] Narumi Y, Aoki Y, Niihori T, Neri G, Cavé H, Verloes A, et al. Molecular and clinical characterization of cardio-facio-cutaneous (CFC) syndrome: Overlapping clinical manifestations with Costello syndrome. *Am J Med Genet Part A* 2007; 143A:799–807.

Based on the oral lesions, presumptive diagnosis of keratotic lichen planus was established. In order to reach definitive diagnosis, a biopsy of the cheek mucosa was taken and the following diagnostic methods were indicated: routine histopathological studies (hematoxylin-eosin and PAS staining), and direct immunofluorescence. Results of the histopathological and immunohistochemical study were consistent with oral lichen planus.

Given the presence of lesions at both commissures of the lips and on the dorsal surface of the tongue associated with Candida infection, said lesions were swabbed and sent for mycological and bacteriological examination, identification of the isolated species, and antibacterial and antifungal susceptibility tests, in order to rule out superadded infection and to orient treatment. The isolated microorganism was Candida albicans, and administration of an antifungal drug was indicated in order to eradicate the infection.

Remission of red atrophic erythematous patches, as well as marked improvement of the verrucous areas were observed at the subsequent follow-up visits after administration of antifungal treatment for Candidiasis and of local corticoids for OLP. Only some areas of white plaques and the presence of white reticular lesions of OLP were observed.

Discussion .3

Within the ectodermal anomalies, follicular hyperkeratosis in the arms, legs, and face are the most frequent cutaneous abnormalities in people with CFC syndrome (8). However, although both tissues derive from the same embryonic layer, no oral mucosa anomalies have been reported to date.

- [7] Roberts AE. The clinical phenotype of cardiofaciocutaneous syndrome; in Zenker M (Ed): Noonan Syndrome and related disorders- A Matter of Deregulated Ras Signaling. Monogr Hum Gent Basel, Karger, 2009, vol 17, pp 66-72.
- [8] Roberts A, Allanson J, Jadico SK, et al. The cardiofaciocutaneous syndrome. *J Med Genet.* 2006; 43(11):833-42.
- [9] Weston G, Payette M. Update on lichen planus and its clinical variants. *Int J Womens Dermatol.* 2015; 1(3):140–149. doi:10.1016/j.ijwd.2015.04.001
- [10] Boyd AS, Neldner KH. Lichen planus. *J. Am. Acad. Dermatol.* 1991 Oct; 25(4):593-619
- [11] Wagner G., Rose C., Sachse M.M. Clinical variants of lichen planus. *J Dtsch Dermatol Ges.* 2013; 11(4):309– 319.
- [12] Gorouhi F, Davari P, Fazel N. Cutaneous and mucosal lichen planus: a comprehensive review of clinical subtypes, risk factors, diagnosis, and prognosis. *ScientificWorldJournal.* 2014; 2014:742826. doi:10.1155/2014/742826.
- [13] Le Cleach L, Chosidow O. Clinical practice. Lichen planus. *N. Engl. J. Med.* 2012 Feb 23; 366(8):723-32
- [14] Aghbari SMH, Abushouk AI, Attia A, Elmaraezy A, Menshawy A, Ahmed MS, Elsaadany BA, Ahmed EM. Malignant transformation of oral lichen planus and oral lichenoid lesions: A meta-analysis of 20095 patient data. *Oral Oncol.* 2017 May; 68:92-102. Doi: 10.1016/j.oraloncology.2017.03.012.
- [15] Richards D1. Malignant transformation rates in Oral Lichen Planus. *Evid Based Dent.* 2018 Dec; 19(4):122. Doi: 10.1038/sj.ebd.6401349.



IJSURP Publishing Academy

International Journal Of Scientific And University Research Publication

Multi-Subject Journal

Editor.

International Journal Of Scientific And University Research Publication



+965 99549511



+90 5374545296



+961 03236496



+44 (0)203 197 6676

www.ijsurp.com